

UNDERWOOD'S BONNETED BAT (*EUMOPS UNDERWOODI*):  
FIRST RECORD IN THE BAJA CALIFORNIA PENINSULA

Patricia Cortés-Calva<sup>1,2</sup>, Sergio Ticul Álvarez-Castañeda<sup>1</sup>,  
Julio M. Hernández-Gutiérrez<sup>1</sup>, and Mayra de la Paz Cuevas<sup>1</sup>

**ABSTRACT.**—This paper reports the first record of *Eumops underwoodi* in the Baja California peninsula. This record expands the species' distributional range in Mexico along the peninsula. We propose 2 plausible biogeographical explanations. First, the species has the ability to travel across the Gulf of California from the coast of Sonora to the eastern side of the Baja California peninsula and there establish. Second, there is a continuous distribution of the species throughout the mountain ranges that run longitudinally along the Baja California peninsula from the United States border down to the southern tip. We collected one male specimen of *E. underwoodi* in xeric-riparian habitat. We suggest a continuous distribution pattern of this species along the cliffs in the Baja California peninsula.

**RESUMEN.**—Se realiza el primer registro de *Eumops underwoodi* en la península de Baja California, con lo que se amplía su intervalo de distribución a lo largo de la península y para México. Se proponen dos explicaciones biogeográficas para este registro. La primera analiza la capacidad de viajar a través del Golfo de California, desde la costa de Sonora hasta la península de Baja California. La segunda, considera que existe una distribución continua de *E. underwoodi* a través de las cadenas montañosas presentes longitudinalmente a lo largo de la península de Baja California, desde la frontera de Estados Unidos hasta el extremo sur. Se colectó un ejemplar macho de *E. underwoodi* en hábitat xérico ripario. Se considera que esta especie tiene un patrón de distribución continuo a lo largo de los acantilados de la península de Baja California.

The genus *Eumops* (family Molossidae) includes 14 species (Eger 2007, McDonough et al. 2008, Baker et al. 2009, Gregorin 2009), 6 of which are distributed in Mexico: *Eumops auripendulus* (Best et al. 2002), *E. nanus* (Eger 2007), *E. ferox* (Baker et al. 2009), *E. hansae* (Best et al. 2001), *E. perotis* (Best et al. 2001), and *E. underwoodi* (Kiser 1995).

The distribution of the genus *Eumops* comprises a variety of habitats—coastal, xeric, mesic, and forest (Kiser 1995, Best et al. 1996, 2001, 2002, Hunt et al. 2003)—occurring at elevations ranging from sea level to 3000 m (Eger 1977). Specimens of *Eumops*, and other bats associated with xeric habitats, have been collected over artificial water bodies (Cockrum and Gardner 1960, Barbour and Davis 1969).

Two *Eumops* species, *E. perotis* and *E. underwoodi*, have been recorded in the states of Sinaloa and Sonora on the eastern side of the Gulf of California basin (Hall 1981, Arroyo-Cabrales 1999). *Eumops perotis* was recently recorded in Sierra de Juárez, Baja California (Martínez-Gallardo et al. 2008), which is the species' southernmost recorded location. The

distribution of *E. underwoodi* stretches from Arizona to Nicaragua (Hall 1981, Kiser 1995, Sánchez and Romero 1995, Medellín et al. 1997, Mac-Swiney González et al. 2003, Simmons 2005).

We sampled for bats in the “La Ascension” wetland located 18 km N and 19 km W of San Bruno, Baja California Sur (26°23'14.95" N, 111°36'3.51" W; Fig. 1), on 13 July 2010. We placed 2 mist nets (8 × 2.5 m) over the water for 5 hours and monitored them regularly. The water body is a permanent spring with a depth ranging between 20 and 40 cm and with the presence of small fish and some beetles. The vegetation near the collection site is dominated by date palm tree (*Phoenix dactylifera*), tree tobacco (*Nicotiana glauca*), cattail (*Typha* sp.), and palo blanco (*Lysiloma candidum*), with an average height of 240 cm, as well as apan (*Apan bebbia*), eucnide (*Eucnide* sp.), yellow orchid vine (*Callaeum macropterum*), and heliotropium (*Heliotropium curassavium*), with an average height of 150 cm. The surrounding vegetation consists of thorn scrub including cacti (*Pachycereus pringlei*), mesquite (*Prosopis*

<sup>1</sup>Centro de Investigaciones Biológicas del Noroeste (CIBNOR), S.C. Instituto Politécnico Nacional 195, La Paz, Baja California Sur 23096, México.

<sup>2</sup>E-mail: pcortes04@cibnor.mx




Fig. 1. Map of the Baja California Peninsula showing the collection site (star).

*palmeri*), cholla (*Cylindropuntia cholla*), Palo Verde (*Cercidium microphyllum*), limberbush (*Jatropha cuneata*), and sweet acacia (*Acacia farnesiana*), with an average height of 250 cm.

We collected one *E. underwoodi* male (CIB catalog number 18139) on the xeric-riparian habitat along the watercourses. It was captured at 23:40 (4 hours after sunset) on the second line (top) of the net placed on the river bed. The scrotum was not evident, but the throat gland was evident and gave off a slightly musky odor. In the laboratory, we recorded morphometric measurements of the skull and established the pelage color according to the Munsell color system (Munsell 2000).

External measurements (mm) of the specimen were as follows: total length 153.0, tail 55.0, hindfoot 12.0, ear 26.0, and forearm 66.0; body mass was 44.0 g. Cranial measurements (mm) for the specimen were as follows: greatest length of skull 28.30, condylobasal length 26.30, zygomatic breadth 17.30, mastoid breadth 13.46, braincase breadth 14.45, braincase depth 11.50, palatal length 11.22, breadth across upper molars 11.13, maxillary tooth row length 11.66, width across upper canines 6.19, postorbital constriction 5.60, mandible length 21.20, and mandibular tooth row length 12.60.

The dorsal color was a grayish brown (10YR5/2) with a white hair base (10YR8/1). The ventral color was light gray (10YR7/2). The dorsal hair was soft and short compared to hair on the hindquarters, which was longer (approximately 12.8 mm). The ears were long and when pushed forward reached the tip of the nose. The specimen had short hairs on the edge of the calcaneus.

The *E. underwoodi* specimen and tissue were deposited in the Centro de Investigaciones Biológicas del Noroeste (CIB Mammal Collection), as per the recommendations of the Animal Care and Use Committee of the American Society of Mammalogists (1998).

Other bats captured included 3 *Nyctinomops femorosaccus*, 1 *Mormoops megalophylla*, 4 *Leptonycteris yerbabuena*, 22 *Antrozous pallidus*, 1 *Corynorhinus townsendii*, 4 *Eptesicus fuscus*, 4 *Myotis volans*, and 6 *Parastrellus hesperus*. These 8 species represent 4 families: Molossidae, Mormoopidae, Phyllostomidae, and Vespertilionidae. The first species collected were *Parastrellus* and *Myotis* (1 hour after sunset), followed by *Nyctinomops*, *Corynorhinus*, and *Leptonycteris* (2 hours after sunset). From 22:30 (3 hours after sunset) the most abundant species were *Antrozous pallidus* and *Parastrellus*

*hesperus*. Capture success was 6 bats per m<sup>2</sup> of net per hour per night.

*Eumops underwoodi* is widely distributed from Arizona (USA) to Nicaragua (Simmons 2005). The specimen we collected is the only known record in the Baja California peninsula (Villa-R. 1967, Hall 1981, Arroyo-Cabrales 1999). Our capture of *E. underwoodi* is biogeographically important because the closest known record is 1 km W of San José de Guaymas, Sonora (Hall 1981), approximately 175 km across the Gulf of California. According to the original description (Benson 1947) and systematics studies made by Eger (1977), the specimen we collected corresponds to *E. underwoodi sonoriensis*. This subspecies is smaller (body, forearm, and skull) than *E. u. underwoodi* and shows some differences in pelage color, in addition to its allopatric distribution (Gregorin 2009). The specimen collected in the Baja California peninsula had a coloration slightly paler than *E. u. underwoodi* and similar to Benson's (1947) original description of pelage color (brown). It had a small skull with measurements near the minimums given by Benson (1947).

We propose 2 explanations for the presence of *E. underwoodi* in the Baja California peninsula. First, this species presumably is capable of traveling the 150–200-km width across the Gulf of California from the coast of Sonora to the eastern side of the Baja California peninsula. Some bat species migrate and travel great distances in the breeding season (Ceballos et al. 1997, Hedenström 2009), and some molossid migrate long distances (Cockrum 1969).

The second explanation, and in our opinion the most parsimonious one, proposes a continuous distribution of *E. underwoodi* through the Sierra de Juárez, Sierra San Pedro Mártir, Sierra Guadalupe, and Sierra de la Giganta, which run longitudinally along the Baja California peninsula from the U.S. border to the southern tip. Recent records of congener *E. perotis* on the Baja California peninsula (Martinez-Gallardo et al. 2008) lend supporting evidence to the idea of a continuous distribution of *E. underwoodi* through the peninsula. We also suggest that the heterogeneity that characterizes the Baja California desert environment (Riddle and Hafner 2006) favors the occurrence of *Eumops* (and other bat genera) in this region, given the ecological plasticity of this genus in selecting roosts, foraging place,

and food availability (Fenton 1989, 1990, Siemers et al. 2001, Pio et al. 2010). Slopes and physiography in these mountain ranges are optimal for initiating flight into the open spaces high above the ground where molossid bats typically hunt (Iníguez-Dávalos 2005). Also, vegetation types and structure are present for roosting and for feeding on arthropods, as well as nectar, pollen, and flowers (Cockrum 1960, Schnitzler and Kalko 2001, Tibbitts et al. 2002). More specifically, *Eumops underwoodi* is associated with mountain cliffs and possibly uses cactus cavities as roost sites, as reported for *Eumops* in Sonoyta where this species occupied cavities in saguaro cacti (Tibbitts et al. 2002).

Also, our data on bat species richness and total number of bats on “La Ascension” indicate that this place on the Baja Peninsula is a “hot spot,” with resources (roosts, foraging places, and food availability) important in supporting bat fauna. These wetland environments could provide vegetation types optimal for refuges or roosting.

In conclusion, the biogeographic implications of the record we present herein are critical because *E. underwoodi* likely has a wider distribution throughout the Baja California peninsula than the few records indicate. *Eumops underwoodi* is a large and conspicuous species, but one that is difficult to collect due to its habits of using cliffs for refuge and flying at high altitudes above the substrate (Schmidly 1991). Additionally, specific natural history studies in the Baja California peninsula have not focused on bats, and monitoring and collection efforts are scant. This first record confirms the importance of wetlands in desert environments, since *E. underwoodi* has been reported as having a wet-tropical distribution (Arroyo-Cabrales 1999).

This project was undertaken with a grant provided by the *Comisión Nacional para el Conocimiento y Uso de la Biodiversidad* (CONABIO) [GT033 “Inventario de los murciélagos de las zonas desérticas y semidesérticas del norte de México” to Patricia Cortés-Calva]. Scientific collection licenses were granted by SEMARNAT (FAUT-0051). We acknowledge D. Dorantes for the English edition.

#### LITERATURE CITED

ANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF MAMMALOGISTS. 1998. Guidelines for the

- capture, handling, and care of mammals as approved by the American Society of Mammalogists. *Journal of Mammalogy* 79:1416–1431.
- ARROYO-CABRALES, J. 1999. Familia Molossidae. Pages 183–198 in S.T. Álvarez-Castañeda and J.L. Patton, editors, *Los mamíferos del noroeste de México*. Centro de Investigaciones Biológicas del Noroeste, SC., México.
- BAKER, R.J., M.M. McDONOUGH, V.I. SWIER, P.A. LARSEN, J.P. CARRERA, AND L.K. AMMERMAN. 2009. New species of bonneted bat, genus *Eumops* (Chiroptera: Molossidae) from the lowlands of western Ecuador and Peru. *Acta Chiropterologica* 11:1–13.
- BARBOUR, R.W., AND W.H. DAVIS. 1969. *Bats of America*. University Press of Kentucky, Lexington, KY.
- BENSON, S. 1947. Description of a mastiff bat (Genus *Eumops*) from Sonora, Mexico. *Proceedings of the Biological Society of Washington* 60:133–134.
- BEST, T.L., J.L. HUNT, L.A. MCWILLIAMS, AND K.G. SMITH. 2001. *Eumops hansae*. *Mammalian Species* 687:1–3.
- \_\_\_\_\_. 2002. *Eumops auripendulus*. *Mammalian Species* 708:1–5.
- BEST, T.L., W.M. KISER, AND P.F. FREEMAN. 1996. *Eumops perotis*. *Mammalian Species* 534:1–8.
- CEBALLOS, G., T.H. FLEMING, C. CHAVEZ, AND J. NASSAR. 1997. Population dynamics of *Leptonycteris curasoae* (Chiroptera: Phyllostomidae) in Jalisco, Mexico. *Journal of Mammalogy* 78:1220–1230.
- COCKRUM, E.L. 1960. Distribution, habitat and habits of the mastiff bat, *Eumops perotis*, in North America. *Journal of the Arizona Academy of Science* 1:79–84.
- \_\_\_\_\_. 1969. Migration in the guano bat, *Tadarida brasiliensis*. *University of Kansas Museum of Natural History Miscellaneous Publication* 51:303–336.
- COCKRUM, E.L., AND A.L. GARDNER. 1960. Underwood's mastiff bat in Arizona. *Journal of Mammalogy* 41:510–511.
- EGER, J.L. 1977. Systematics of the genus *Eumops* (Chiroptera: Molossidae). *Life Sciences Contributions, Royal Ontario Museum* 110:1–69.
- \_\_\_\_\_. 2007. Family Molossidae P Gervais, 1856. Pages 399–440 in A.L. Gardner, editor, *Mammals of South America*. Volume 1, Marsupials, xenarthrans, shrews, and bats. University of Chicago Press, Chicago, IL.
- FENTON, M.B. 1989. Head size and the foraging behaviour of animal-eating bats. *Canadian Journal of Zoology* 67:2029–2035.
- \_\_\_\_\_. 1990. The foraging behaviour and ecology of animal eating bats. *Canadian Journal of Zoology* 68:411–422.
- GREGORIN, R. 2009. Phylogeny of *Eumops* Miller, 1906 (Chiroptera: Molossidae) using morphological data. *Acta Chiropterologica* 11:247–258.
- HALL, E.R. 1981. *The mammals of North America*. John Wiley & Sons, New York, NY.
- HEDENSTRÖM, A. 2009. Optimal migration strategies in bats. *Journal of Mammalogy* 90:1298–1309.
- HUNT, J.L., L.A. MCWILLIAMS, T.L. BEST, AND K.G. SMITH. 2003. *Eumops bonariensis*. *Mammalian Species* 733:1–5.
- ÍÑIGUEZ-DÁVALOS, L.I. 2005. *Eumops underwoodi*. Pages 321–322 in G. Ceballos and G. Oliva, editors, *Los Mamíferos Silvestres de México*. Fondo de Cultura Económica, Comisión Nacional para el Conocimiento y Uso de la Biodiversidad, México, D.F.
- KISER, W.M. 1995. *Eumops underwoodi*. *Mammalian Species* 516:1–4.
- MAC-SWINEY GONZÁLEZ, M.C., J. SOSA-ESCALANTE, AND C.I. SELEM-SALAS. 2003. Ampliación en la distribución de *Eumops underwoodi* Goodwin, 1940 (Chiroptera: Molossidae) en la Península de Yucatán, México. *Revista Mexicana de Mastozoología* 7:55–57.
- MARTÍNEZ-GALLARDO, R., A.A. GUEVARA CARRIZALES, R. COUOH DE LA GARZA, AND E. FLORES ROJAS. 2008. Primer registro de *Eumops perotis* (Chiroptera: Molossidae) en la península de Baja California. *Universidad y Ciencia* 24:189–190.
- McDONOUGH, M.M., L.K. AMMERMAN, R.M. TIMM, H.H. GENOWAYS, P.A. LARSEN, AND R.J. BAKER. 2008. Speciation within bonneted bats (genus *Eumops*): the complexity of morphological, mitochondrial, and nuclear data sets in systematics. *Journal of Mammalogy* 89:1306–1315.
- MEDELLÍN, R.A., H.T. ARITA, AND O. SÁNCHEZ. 1997. Identificación de los murciélagos de México. Clave de campo. *Publicaciones Especiales, Asociación Mexicana de Mastozoología, A.C.*, México.
- MUNSELL. 2000. *Munsell soil color charts*. GretagMacbeth. New Windsor, New York, NY.
- PIO, D.V.V., F. CLARKE, M.I. MACKIE, AND P.A. RACEY. 2010. Echolocation calls of the bats of Trinidad, West Indies: is guild membership reflected in echolocation signal design? *Acta Chiropterologica* 12:217–229.
- RIDDLE, B.R., AND D. HAFNER. 2006. Biogeografía histórica de los desiertos cálidos de Norteamérica. Pages 57–65 in E. Vázquez-Domínguez and D.J. Hafner, editors, *Genética y mamíferos mexicanos: presente y futuro*. New Mexico Museum of Natural History and Science Bulletin, New Mexico.
- SÁNCHEZ, H.C., AND L. ROMERO. 1995. Murciélagos de Tabasco y Campeche. Una propuesta para su conservación. Cuadernos 24. Instituto de Biología de la Universidad Nacional Autónoma de México, México.
- SCHMIDLY, D.J. 1991. *The bats of Texas*. Texas A&M University Press, College Station, TX.
- SCHNITZLER, H.U., AND E.K.V. KALKO. 2001. Echolocation in insect-eating bats. *BioScience* 51:557–569.
- SIEMERS, B.M., E.K.V. KALKO, AND H.-U. SCHNITZLER. 2001. Echolocation and signal plasticity in the Neotropical bat *Myotis nigricans* (Schinz, 1821) (Vespertilionidae): a convergent case with European species of *Pipistrellus*? *Behavioural Ecology and Sociobiology* 50:317–328.
- SIMMONS, N.B. 2005. Order Chiroptera. Pages 312–529 in D.E. Wilson and D.M. Reeder, editors, *Mammal species of the world: a taxonomic and geographic reference*. 3rd edition. Johns Hopkins University Press, Baltimore, MD.
- TIBBITTS, T., A. PATE, Y. PETRYSZYN, AND B. BARNS. 2002. Determining foraging and roosting areas for Underwood's mastiff bat (*Eumops underwoodi*) using radio-telemetry, at Organ Pipe Cactus National Monument, Arizona. Final Summary Report, Year 2 – December 2002. Appendix G, Ecological Monitoring Program Report, 1997–2005. Organ Pipe Cactus National Monument, Ajo, AZ. 14 pp.
- VILLA-R., B. 1967. Los murciélagos de México. Su importancia en la economía y la salubridad – su clasificación sistemática. Instituto de Biología, Universidad Nacional Autónoma de México, D.F.

Received 21 September 2011

Accepted 25 April 2012