



---

4-30-2004

### Biogeographic and taxonomic relationships among the mountain snails (Gastropoda: Oreohelicidae) of the central Great Basin

Mark A. Ports

*Great Basin College, Elko, Nevada*

Follow this and additional works at: <https://scholarsarchive.byu.edu/wnan>

---

#### Recommended Citation

Ports, Mark A. (2004) "Biogeographic and taxonomic relationships among the mountain snails (Gastropoda: Oreohelicidae) of the central Great Basin," *Western North American Naturalist*. Vol. 64 : No. 2 , Article 1.

Available at: <https://scholarsarchive.byu.edu/wnan/vol64/iss2/1>

This Article is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in Western North American Naturalist by an authorized editor of BYU ScholarsArchive. For more information, please contact [scholarsarchive@byu.edu](mailto:scholarsarchive@byu.edu), [ellen\\_amatangelo@byu.edu](mailto:ellen_amatangelo@byu.edu).

BIOGEOGRAPHIC AND TAXONOMIC RELATIONSHIPS  
AMONG THE MOUNTAIN SNAILS (GASTROPODA: OREOHELICIDAE)  
OF THE CENTRAL GREAT BASIN

Mark A. Ports<sup>1</sup>

ABSTRACT.—Described here are 4 species of mountain snails, *Oreohelix*, isolated on mountains in the central Great Basin of Nevada and Utah since the end of the Pleistocene. Forty-three mountains were searched during an 18-year period, resulting in 24 mountains found with no oreohelicids present. One population, *Oreohelix loisae* (19 mm to 23 mm in shell diameter), is described here as a new species related to, but geographically isolated from, the species *Oreohelix nevadensis* (17 mm to 22 mm diameter). *Oreohelix loisae* is present only in the Goshute Mountains while *O. nevadensis* is represented in 3 geographically adjacent ranges in the central Great Basin. These 2 species are possibly related to the *Oreohelix haydeni* group from the northern Wasatch Range. The subspecies *Oreohelix strigosa depressa* (15 mm to 21 mm diameter) is present on 11 ranges from western Utah west to east central Nevada. This subspecies is closely related to populations found today in the northern Wasatch Mountains of Utah. The smallest species in diameter (8 mm to 14 mm), *Oreohelix hemphilli*, is centered in the central Great Basin and found on 16 ranges often in sympatry with 1 or 2 of the larger conspecifics. Both qualitative and quantitative information on shell characters and soft anatomy is provided here for these 4 species. Shell characters, soft anatomy, geographical isolation, and statistical analysis suggest that 4 distinct species inhabit the central Great Basin today. Xeric and calciphilic species include *O. hemphilli* and *O. loisae*, while *O. strigosa* and *O. nevadensis* typically are associated with permanent water and both metamorphic and limestone mountains.

*Key words:* mountain snails, *Oreohelix*, Great Basin biogeography, taxonomy, *Oreohelix loisae* sp. nov.

The genus *Oreohelix* (Stylommatophora: Oreohelicidae) is widely distributed but localized throughout the Rocky Mountains, Great Basin, and southwestern regions of North America (Pilsbry 1939). Recent work (Solem 1975, Frest and Johannes 1997) has identified 2 centers of diversity in this genus, the 1st from the Salmon River drainage of Idaho and the 2nd from the northern Wasatch Mountains of Utah and Bear River Range of southern Idaho (Pilsbry 1939, Clarke and Hovingh 1991).

Pilsbry (1939) described 3 species of *Oreohelix* from the central and eastern Great Basin mountain ranges of Nevada: *O. hemphilli* Newcomb, 1869, in the White Pine Range, *O. nevadensis* Berry, 1932, in the Schell Creek Range, and *O. strigosa* Gould, 1846 (subspecies *depressa* Cockerell 1890), from the Snake Range. Roscoe (1954) listed *O. subrudis* Reeve, 1854, and *O. eurekaensis* J. Henderson and Daniels, 1916, from the Deep Creek Range in western Utah. During visits in 1996 and 1997, however, the author found a medium-sized shell of *O. strigosa* and smaller-sized shell of *O. hem-*

*philli* in the Deep Creek Range. Apparently, Roscoe (1954) misidentified these 2 species possibly due to the wide variation in shell morphology in both species as they occur throughout Utah, Colorado, and Idaho (Brandauer 1988, Clarke and Hovingh 1991, Frest and Johannes 1997).

Fieldwork in the Great Basin, begun in 1982 by Pratt (1985) and continued by Ports (1986–2002), has increased knowledge of the biogeography of the genus *Oreohelix* in the central Great Basin and suggests that this region may be a 3rd center of oreoheliced diversity as predicted by Bequaert and Miller (1973). Today these populations are fragmented and isolated on mountain ranges throughout the region with little opportunity for gene flow and dispersal across desert valleys, similar to other faunal groups such as high-elevation-adapted small mammals in the Great Basin (Brown 1971).

In this paper I describe the biogeography, shell and soft anatomy, and ecological variation in 3 little-known species of *Oreohelix* as

<sup>1</sup>Biology Department, Great Basin College, 1500 College Parkway, Elko, NV 89801.

listed for Nevada and present as new to science a previously unknown oreohelcid from the Goshute Mountains on the Nevada–Utah border in the north central Great Basin. This new species is related to the *Oreohelix haydeni* group from the Wasatch Mountains of Utah.

#### METHODS AND MATERIALS

From 1986 to 2002, I sought land snails in 43 mountains throughout Nevada and eastern Utah. Shells were organized and stored by lots and stations in a private collection at Great Basin College. Latitude and longitude of each station were determined using U.S. Geological Survey topographic maps and confirmed using a Garmin GPS II or III.

At each station empty shells of all age classes were collected by hand and by digging into rock slides using a rock hammer. Eight to 20 live specimens were collected from each station. These specimens were relaxed in menthol for 12 hours and killed by immersing in boiling water for 30 seconds. Bodies were pulled from their shells, preserved in 75% ethanol for 24 hours, fixed in 5% formalin for 24 hours, and transferred to 50% ethanol for dissection purposes (Frest and Johannes 1997).

Dissection of soft tissue was carried out under a Leicca 40–60X dissection scope. Genitalia were measured in millimeters using vernier digital calipers and drawn from preserved specimens with pen and ink. The radula was prepared and studied on temporary slides using a 1% analine blue dye under a compound light microscope. Empty shells were scanned using Hewlett Packard Precision ScanPro 1.01 in apical, basal, and aperture views, following which images were saved and then printed. The taxonomic description of the new species is according to recommendations of Winston (1999). The following abbreviations were used for repository institutions: ANSP, Academy of Natural Sciences, Philadelphia, for the holotype and paratypes of *O. loisae*, topotypes of *O. nevadensis*, and *O. strigosa*; SBMNH, Santa Barbara Museum of Natural History, paratypes of *O. loisae* and topotypes of *O. nevadensis*, *O. strigosa depressa*, and *O. hemphilli*; MAP, Mark A. Ports catalogue numbers. Catalogue numbers, localities, species, and dates of collection are listed in the appendix. A Mann-Whitney U-test was used to compare morphometric data between *O. strigosa*, *O. nevadensis*, and *O. loisae*.

#### RESULTS

##### Taxonomy and Biogeography of *Oreohelix loisae*

The oreohelcid herein is described as a new species: Order Stylommatophora; Family Oreohelicidae; Genus *Oreohelix* Pilsbry, 1904; *Oreohelix loisae* M.A. Ports, sp. nov. Holotype: ANSP 407556 (Figs. 1, 2, Table 1). Paratypes: ANSP 408362, SBMNH 345735.

DIAGNOSIS.—Description of shell characteristics: Holotype (Fig. 1, 3 views of the holotype): A large shell compared with other Great Basin oreohelicids (holotype, 19.1 mm diameter and 11.4 mm height) with 5.5 whorls. Shell depressed, broadly conical, spire forming an angle of 100 degrees. Shell convex both on apical and basal surfaces with a strongly rounded peripheral body whorl. Yellow-brown periostracum over the apical surface, lighter on the basal surface. Three narrow, yellow-brown bands with the peripheral band above the aperture which descends in front. No banding below. An ovate-lunate aperture, 9.4 mm in diameter and 8.5 mm in height (Table 1). Thin outer lip not deflected, parietal callus thick and reflected slightly over the umbilicus. Umbilicus deep and moderately wide, 4.6 mm in diameter, making up 23% of the shell diameter. Four faint to obsolete beaded lirae on the basal surface and 4 faint beaded lirae on the apical surface. Fine to moderately coarse, oblique striae on body whorls. Faint periostracal wreaths associated with the beaded lirae, but no periostracal fringes on the peripheral whorl. Whorl sutures moderately impressed between penultimate and body whorls, aperture descends below the body whorl.

Paratype specimens vary from the holotype in shell size (17.7–21.8 mm shell diameter; 0.59–0.61 ratio of height/diameter; Table 1), shell color (a light caulk), lack of bands or the presence of banding, distinct angular whorls to rounded body whorls, and degree of beaded lirae on the basal and apical surfaces. Juvenile shells are strongly angular, including aperture whorl, 4 fine and distinct complete lirae on both the apical and basal surface; juveniles have a much more depressed spire than adult shells with a large, ovate-lunate aperture. Periostracum in juveniles is dark brown, a color that accentuates the fine lirae, 3 distinct narrow brown bands, and shallow sutures, umbilicus is narrow, 3.5–whorls. The mean radula formula



Fig. 1. Apical, aperture, and basal view of the holotype specimen of *Oreohelix loisae*, sp. nov. Shell diameter = 21 mm; shell height = 13 mm.




Fig. 2. Reproductive system of *Oreohelix loisae* Ports, sp. nov. Holotype from Christmas Tree Canyon, NV (ANSP 407556). Scale bar = 1 mm. Key: ag = albumin gland, ep = epiphallus, fo = free oviduct, prm = penis retractor muscle, ps = penis sack, sp = spermatheca, ut = uterus, va = vagina, vd = vas deferens. Talon is located on median side of the albumen gland.

for the holotype and 15 dissected paratypes is 24-1-24 (Table 1). Rows of teeth per radula ranged from 10 to 150, the central tooth is a rounded unicuspid on a tall broad plate with no ectocones; lateral teeth 1-15, have a pronounced lateral ectocone and a sharp pointed cuspid; marginal teeth 16-22, degenerate from distinctly bicuspid with small, lateral ectocones to very small unicuspid marginal teeth with no ectocones, 23-24.

DESCRIPTION OF GENITALIA OF HOLOTYPE (FIG. 2).—The penis sac is relatively long (9.98 mm) and narrow throughout and is twisted into a tight S shape at the distal end. The distal end of the penis sac is flattened laterally and is the site of attachment for both the ribbonlike penis retractor muscle and the narrow epiphallus. Internally, 35% of the proximal end of the penis sac is lined with 4-5 regular, longitudinal pilasters, while the remaining internal surface is lined with dense papillae. The epiphallus is relatively long (5.17 mm) and slender, tapering to a thin attachment at both the distal and proximal ends. The distal end of the vas deferens runs along the side of the vagina, coiling around the base of the vagina to enter the apex of the epiphallus in the medial surface between the penis sac and the vagina. The vagina is narrow at the base and broadens at the junction of the free oviduct and the uterus. The spermathecal sac is relatively large and oval in shape and the albumin gland is short and rounded with a sharply recurved black talon. The hermaphroditic duct is highly coiled and compressed and the ovotestes are relatively short and dense.

The genitalia of 25 dissected paratypes varied in having a penis sac length of 10.1 to 12.9 mm, internal ribbed portion of penis sac of 3.79 mm to 4.39 mm in length, a ratio of penis sac length/shell diameter of 0.53 to 0.67, and a ratio of internal ribbed portion/total penis sac length of 0.34 to 0.38 (Table 1).

TYPE MATERIAL.—Holotype: ANSP 407556; Christmas Tree Canyon, Goshute Mountains, Elko County, NEVADA, 40 25.566'N and -144 16.233'W. 2784 m. Paratypes: SBMNH 345735; Felt Wash, Goshute Mountains, Elko County, NEVADA, 40 27.347'N and -114 16.105'W. 2477 m (Fig. 5, number 16).

TABLE 1. Measurements of shell characters, soft anatomy, and ratios of 14 populations of *Oreohelix* from the central Great Basin of North America. Measurements represent mean values in millimeters for shell diameter, umbilicus width, and penis length.

Species	Shell diameter	Shell height/shell diameter	Umbilicus width	Radula formula	Penis length	Penis length/shell diameter	Ribbed portion/penis length
<i>O. nevadensis</i> Schell Creek Range, <i>n</i> = 16	19.9	0.64	4.63	28-1-28	13.3	0.77	0.31
<i>O. loisae</i> Goshute Mountains, <i>n</i> = 10	19	0.59	4.91	24-1-24	10.1	0.53	0.38
<i>O. loisae</i> Goshute Mountains, <i>n</i> = 6	19	0.61	4.4	24-1-24	12.9	0.67	0.34
<i>O. strigosa depressa</i> Ruby Mountains, <i>n</i> = 11	15.2	0.66	3.35	30-1-30	10	0.6	0.35
<i>O. strigosa depressa</i> Ruby Mountains, <i>n</i> = 13	16.3	0.68	3.48	30-1-30	10.8	0.66	0.37
<i>O. strigosa depressa</i> Pequop Mountains, <i>n</i> = 8	16.5	0.61	3.54	30-1-30	10.3	0.62	0.35
<i>O. strigosa depressa</i> Snake Range, <i>n</i> = 10	17.6	0.63	4.24	30-1-30	14.1	0.8	0.38
<i>O. strigosa depressa</i> Schell Creek Range, <i>n</i> = 8	16.9	0.67	3.66	30-1-30	9.9	0.59	0.327
<i>O. strigosa depressa</i> Pilot Range, <i>n</i> = 8	18.2	0.70	3.72	30-1-30	7.7	0.42	0.37
<i>O. strigosa depressa</i> Deep Creek Range, <i>n</i> = 8	18.4	0.64	4.46	30-1-30	9.9	0.54	0.38
<i>O. strigosa depressa</i> Egan Range, <i>n</i> = 6	16.2	0.64	4.56	30-1-30	10.6	0.66	0.37
<i>O. hemphilli</i> White Pine Range, <i>n</i> = 12	11.4	0.6	2.4	22-1-22	4	0.36	0.38
<i>O. hemphilli</i> Snake Range, <i>n</i> = 6	12.5	0.64	3.1	24-1-24	4	0.38	0.6
<i>O. hemphilli</i> Ruby Mountains, <i>n</i> = 8	8.0	0.61	2.6	22-1-22	2.6	0.32	0.36

ETYMOLOGY.—This species is named in honor of Lois K. Ports, my field companion, friend, and wife.

Taxonomy and Biogeography  
of *Oreohelix nevadensis*  
S.S. Berry, 1932

The Schell Creek mountainsnail (*O. nevadensis*; SBMNH 345735) is the largest species of this genus in the Great Basin (mean diameter of 19.9 mm) and according to Pilsbry (1939) is most similar to the diverse *Oreohelix haydeni* (Gabb, 1869) group from north central Utah in the Wasatch Mountains. The shell of *O. nevadensis* (Fig. 3) has weak "spiral striae" or beaded lirae and broad, distinctive bands against a

dark brown periostracum. Juvenile specimens have an angular peripheral whorl and flattened spire. Mean shell diameter is 19.9 mm, the umbilicus is deep and wide at 4.63 mm, shell height/shell diameter ratio is 0.64, and the species possesses a mean radula formula of 28-1-28 (Table 1). The genitalia (Fig. 4) of this snail are similar in size (mean 12.9 mm penis sac length) and shape to that of *O. loisae*. Although the penis sac length/shell diameter ratio is higher than all other species examined, the ribbed portion/penis sac length is smallest, due to the large shell size and the relatively large ribbed portion length of 4.02 mm (Table 1). Comparisons of shell diameter ( $P < 0.001$ ), penis length ( $P < 0.001$ ), and the ratio of penis




Fig. 3. Apical, aperture, and basal view of *Oreohelix nevadensis*. Shell diameter = 23 mm; shell height = 15 mm.




Fig. 4. Reproductive system of *Oreohelix nevadensis* S.S. Berry, 1932. Topotype from Cleve Creek, Schell Creek Range, NV (ANSP 4018885). Scale bar = 1 mm. Key: ag = albumin gland, ep = epiphallus, fo = free oviduct, prm = penis retractor muscle, ps = penis, sp = spermatheca, ta = talon, ut = uterus, va = vagina, vd = vas deferens.

sac length/shell diameter were significantly different ( $P < 0.001$  level) for *O. loisae* and *O. nevadensis*. Mean penis sac length and shell diameter (Table 1) separate this species from the others examined, but they are closest to *O. loisae*.

The specimens of *O. nevadensis* examined (ANSP 401885, SBMNH 345735; Appendix), come from the type locality on Cleve Creek (Pilsbry 1939) on the east slope of the Schell Creek Range and from 2 stations on the west slope. I also collected specimens on Goshute Creek and McDermid Creek in the Cherry Creek Range and on Smith Creek in the northern Snake Range (Fig. 5).

This species is limited in its distribution, typically found in small colonies and restricted to the White Pine County and southern Elko County region of Nevada.

Taxonomy and Biogeography  
of *Oreohelix strigosa depressa*  
Cockerell, 1890

The Rocky mountainsnail, *Oreohelix strigosa* (subspecies *depressa*), in the Great Basin has a smaller shell diameter, 4–5 mm less than the same form found in the Wasatch Range of Utah (Pilsbry 1939), and is intermediate in size between the large-shelled *O. nevadensis* and the small-shelled *O. hemphilli* (Table 1). Six of 7 populations examined are intermediate in penis sac length/shell diameter ratio to the other 3 species (Table 1). The Baker Creek sample (Pilsbry 1939) has a significantly larger penis sac length ( $P < 0.001$ ) and a larger penis sac length/shell diameter ratio ( $P < 0.001$  level) when compared with other populations of this species from the Great Basin.

The Lutts Creek population from the Ruby Mountains (Ports 1993; ANSP 401886) had a significantly smaller shell diameter ( $P < 0.001$ ), penis sac length ( $P < 0.001$ ), and penis sac length/shell diameter ( $P < 0.001$ ) compared with *O. loisae* (Table 1) from the Goshute Mountains. Eight populations examined had a radula formula of 30-1-30, the same radula formula found by Jones (1940) in the Wasatch Mountains. The Lutts Creek snail has 2 to 3 distinct, brown apical bands and a periostracum exhibiting a wide range of color variation from caulk white to dark brown (Fig. 6).

This species was located in 11 of 43 mountain ranges sampled in the central Great Basin (Appendix). Populations exist today from the




Fig. 5. Distribution of *Oreohelix* populations in the central Great Basin of North America:

- |                       |                         |                         |
|-----------------------|-------------------------|-------------------------|
| (1) Black Rock Range  | (16) Pequoop Mts.       | (31) Monitor Range      |
| (2) Pine Forest Range | (17) East Humbolt Range | (32) Hot Creek Range    |
| (3) Santa Rosa Range  | (18) Goshute Mts.       | (33) Roberts Mts.       |
| (4) Tuscarora Mts.    | (19) Spruce Mts.        | (34) Diamond Mts.       |
| (5) Snowstorm Mts.    | (20) Cherry Creek Mts.  | (35) Pancake Range      |
| (6) Sheep Creek Mts.  | (21) Ruby Mts.          | (36) White Pine Range   |
| (7) Independence Mts. | (22) Shoshone Range     | (37) Grant Range        |
| (8) Bull Run Mts.     | (23) Pah Rah Range      | (38) Egan Range         |
| (9) Jarbidge Mts.     | (24) Pine Nut Mts.      | (39) Schell Creek Range |
| (10) Good Creek Mts.  | (25) Sweetwater Mts.    | (40) Snake Range        |
| (11) Raft River Range | (26) White Mts.         | (41) Wilson Creek Range |
| (12) Oquirrh Mts.     | (27) Stillwater Range   | (42) Clover Mts.        |
| (13) Stansbury Mts.   | (28) Clan Alpine Mts.   | (43) Pine Valley Mts.   |
| (14) Pilot Range      | (29) Toiyabe Range      | (44) Deep Creek Range   |
| (15) Toana Range      | (30) Toquima Range      | (45) House Range        |




Fig. 6. Apical, aperture, and basal view of *Oreohelix strigosa depressa*. Shell diameter = 18 mm; shell height = 11 mm.




Fig. 7. Apical, aperture, and basal view of *Oreohelix hemphilli*. Shell diameter = 12 mm; shell height = 7 mm.

Pilot Range (ANSP 401887) in the north, to the Egan Range in the south, as far west as the Diamond Range, east to the Deep Creek Range (ANSP 401888), and farther east to the House Range (Appendix, Fig. 5).

#### Taxonomy and Biogeography of *O. hemphilli* (Newcomb, 1869)

The White Pine mountainsnail (SBMNH 345737) has the smallest shell diameter (mean 8–12.5 mm; Table 1) of the Great Basin oreohelicids. Its shell also has distinct whorled lirae on the basal and apical surfaces, and a carinate, keeled, peripheral body whorl (Fig. 7). The short, swollen penis sac, short epiphallus, and stocky vagina (Fig. 8) are most similar to *O. carinifera* from Montana and *O. eurekaensis* from the East Tintic Range (Pilsbry 1939). The type locality from the White Pine Range is intermediate in shell characters and soft anatomy between the 3 populations examined (Table 1). The radula formula for the type locality specimens and the Pearl Creek specimens is 22-1-22, but a radula formula of 24-1-24 was discovered for specimens from Murphy Wash (Table 1).

This species is located in 16 of 43 mountains sampled (Appendix, Fig. 5). The westernmost population is found in central Nevada in the Toiyabe Range, south to the Grant Range, north in the Pilot Range, and east to the Deep Creek Range and House Range of western Utah (Fig. 5). The last 2 localities represent the 1st records for *O. hemphilli* from Utah. In the Mojave Desert 2 very similar species exist (Pilsbry 1939): *O. handi* Pilsbry and Ferriss, 1918, from the Spring Range, and *O. californica* S.S. Berry, 1931, from Clark Mountain.

#### DISCUSSION

At present 4 species of Oreohelicidae exist in the central Great Basin of North America: *Oreohelix loisae*, *O. nevadensis*, *O. strigosa*, and *O. hemphilli*. Of 19 mountain ranges that support oreohelicids, 2 to 3 species exist in sympatry in 11 mountain ranges, within which no species were found to exist below an elevation of 2000 m (Appendix, Fig. 5). Roscoe (1954) reported both *Oreohelix subrudis* and *Oreohelix eurekaensis* from the Deep Creek Range on the Nevada–Utah border. My collections from






Fig. 8. Reproductive system of *Oreohelix hemphilli* Newcomb, 1869. Topotype from Cathedral Canyon, White Pine Range, NV (MAP 994). Scale bar = 1 mm. Key: ag = albumin gland, ep = epiphallus, fo = free oviduct, prm = penis retractor muscle, ps = penis, sp = spermatheca, ta = talon, ut = uterus, va = vagina, vd = vas deferens.

this range suggest that these species are actually *O. strigosa* and *O. hemphilli*, based on shell morphology and genitalia. Both species pairs (*O. strigosa*–*O. subrudis* and *O. hemphilli*–*O. eurekensis*) are very similar in shell morphology and may be synonymous, as Brandauer (1988) suggested for *O. strigosa* and *O. subrudis* in Colorado. *Oreohelix hemphilli* and *O. eurekensis* are very much alike in many shell characters and may be synonymous with *O. eurekensis* in the East Tintic Mountains on the eastern edge of the Great Basin (Pilsbry 1939).

*Oreohelix loisae* is found in only a few dry canyons of the limestone-dominated Goshute Range in north central Great Basin. A large- to medium-sized snail, this species is adapted to limestone rockslides with an understory of *Symphoricarpus oreophilus*, *Rhus trilobata*, *Artemisia tridentata*, and *Holodiscus dumosus*, beneath an overstory of white fir (*Abies concolor*), limber pine (*Pinus flexilis*), and Great Basin bristlecone pine (*Pinus longaeva*).

*Oreohelix loisae* is found from the highest elevation of the range at 2750 m (bristlecone pine and white fir) to 2450 m in the pinyon pine (*Pinus edulis*) and Utah juniper (*Juniperus osteosperma*) woodlands. Except for spring snowmelt and rare summer showers as sources of moisture, there is no open or flowing water in this range. This snail is found in the same rockslides with *O. hemphilli*, but the other 2

species discussed were not detected in this range.

Because of similar shell characters and soft anatomy, I assume that *O. nevadensis* and *O. loisae* are related to, and possibly holophyletic with, *Oreohelix haydeni* from the northern mountains of Utah (Pilsbry 1939). As of this writing, *O. nevadensis* exists in 3 parallel mountains separated by 2 valleys: the North Snake Range on Smith Creek, the Schell Creek Range on Cleve Creek (the type locality; Pilsbry 1939), and on Goshute Creek in the Cherry Creek Range (Appendix, Fig. 5). In each of these ranges, *O. nevadensis* is associated with perennial springs or streams lined with narrow-leaf cottonwood (*Populus angustifolia*), willow (*Salix* sp.), red-osier dogwood (*Cornus stolonifera*), and wild rose (*Rosa woodsii*). At each locality *O. nevadensis* is sympatric with *O. strigosa* in the drainages and *O. hemphilli* on the drier, brush-covered, rocky slopes above the drainage. It is not unusual to find more than 1 species of oreohelicid on a single, isolated mountain range, as shown by Metcalf and Smartt (1997) in New Mexico. Adaptation to microhabitats and dispersal biogeography (Pratt 1985) into the Great Basin mountains may explain why 2 to 3 species coexist in the same mountains. In Logan Canyon of the northern Wasatch Range, I found *O. strigosa* and *O. haydeni* together (Pilsbry 1939), while in the Stansbury Range on the east side of the Great Basin, I found *O. strigosa* and *O. eurekensis* together in the same canyon. Colonization from the east into the Great Basin also may explain why certain species exist where they do today, as suggested for the distribution of montane mammals in the Great Basin by Brown (1971). Pratt (1985) suggests that several species of land snails that have been isolated on mountains in the Great Basin are at risk of extinction.

The most mesic stations in the central Great Basin are occupied primarily by *O. strigosa*. All of the stations listed for this species (Fig. 5, Appendix) are on perennial streams in limestone or metamorphic ranges in woodlands dominated by quaking aspen (*Populus tremuloides*), limber pine, and white fir with an understory of wild rose, red-osier dogwood, and common chokecherry (*Prunus virginiana*). This species of land snail is generally restricted to rock slides and boulders but occasionally is found in forest litter. Similar habitat associations have been described by Henderson (1924)

for oreohelicids at many stations throughout the West.

Much less is known about the biogeography, anatomy, and habitat association of *Oreohelix hemphilli* (Pilsbry 1939). The source of origin may possibly be the East Tintic Mountains, where the similar *Oreohelix eurekensis* exists, or *O. hemphilli* may be endemic to the central Great Basin and has spread east from the central Great Basin. Stations where *O. hemphilli* is found in the Great Basin (Fig. 5, Appendix) are typically xeric and associated with limestone rockslides. An understory of shrubs is usually present, including snowberry, mountain ninebark (*Physocarpus monogynus*), and Rocky Mountain maple (*Acer glabrum*). Overstory trees consist of mixed conifer woodlands of pinyon pine, Utah juniper, and Rocky Mountain juniper (*Juniperus scopularum*) at mid-elevations (2250 m). At higher elevations (2450 m) the conifers consist of bristlecone pine, limber pine, white fir, and Douglas-fir (*Pseudotsuga menziesii*).

More work, including genetic analysis, is currently underway in order to understand the historical biogeography and taxonomic relationships among these 4 species and their relationships with the species of oreohelicids from the eastern edge of the Great Basin and the Wasatch Mountains.

#### ACKNOWLEDGMENTS

The artwork, table, and photographs were provided by Lois K. Ports and Mark E. Ports. I acknowledge the support and facilities used at Great Basin College and appreciate the acceptance and cataloging of snail specimens by the Academy of Natural Sciences of Philadelphia and the Santa Barbara Museum of Natural History. I appreciate comments from Dr. Lawrence E. Stevens and 2 anonymous reviewers. Finally, I thank Lois, Roger, Jacki, and Mark, who endured rough mountain roads and patiently searched through rock slides for snails over many summers.

#### LITERATURE CITED

- BEQUAERT, J.C., AND W.B. MILLER. 1973. The mollusks of the arid Southwest. University of Arizona Press, Tucson. 271 pp.
- BRANDAUER, N.E. 1988. Family Oreohelicidae (Gastropoda: Pulmonata). Colorado University of Colorado Museum, Natural History Inventory of Colorado 9:1-32.
- BROWN, J.H. 1971. Mammals on mountaintops: non-equilibrium insular biogeography. *American Naturalist* 105:467-478.
- CLARKE, A.H., AND P. HOVINGH. 1991. Status survey of selected land and freshwater gastropods in Utah. Draft report, USDI, Fish and Wildlife Service. Eco-search, Inc., Portland, TX. 70 pp.
- FREST, T.J., AND E.J. JOHANNES. 1997. Land snail survey of the lower Salmon River drainage, Idaho. Technical Bulletin 97-18, Bureau of Land Management. Deixis Consultants, Seattle, WA. 142 pp.
- HENDERSON, J. 1924. Mollusca of Colorado, Utah, Montana, Idaho, and Wyoming. University of Colorado Studies 13:65-223.
- JONES, D.T. 1940. A study of the Great Basin land snail, *Oreohelix strigosa depressa*. University of Utah Bulletin 31:1-43.
- METCALF, A.L., AND R.A. SMARTT, EDITORS. 1997. Land snails of New Mexico. New Mexico Museum of Natural History and Science Bulletin 10. 145 pp.
- PILSBRY, H.A. 1939. Land Mollusca of North America (north of Mexico). Volume 1, part 1. Academy of Natural Sciences of Philadelphia Monograph 3(1):1-574.
- PORTS, M.A. 1996. Habitat affinities and distributions of land gastropods from the Ruby Mountains and East Humboldt Range of northeastern Nevada. *Veliger* 39:335-341.
- PRATT, W.L. 1985. Insular biogeography of central Great Basin land snails: extinction without replacement. *Journal of the Arizona-Nevada Academy of Sciences* 20:14-18.
- ROSCOE, T.J. 1954. Some terrestrial gastropods from the Deep Creek Mountains, Juab County, Utah. *Great Basin Naturalist* 14:19.
- SOLEM, A. 1975. Notes on Salmon River oreohelid land snails, with description of *Oreohelix waltoni*. *Veliger* 18:16-30.
- WINSTON, J.E. 1999. Describing species: practical taxonomic procedures for biologists. Columbia University Press, New York. 518 pp.

Received 12 July 2002  
Accepted 27 August 2003

Appendix on the following page.

## APPENDIX

## SPECIMENS EXAMINED

**I. *Oreohelix nevadensis*:** (1) ANSP 401855 (topotype); SBMNH 345735 (topotype), Cleve Creek, east slope of the Schell Creek Range, White Pine County, NEVADA, 39°.247 N, -114°.925 W, 20 May 1994. (2) MAP 832, Goshute Creek, Cherry Creek Range, Elko County, NEVADA, 40°058 N, -115°835 W, 8 August 1988. (3) MAP 807, Smith Creek, North Snake Range, White Pine County, NEVADA, 39°243 N, -114°107 W, 28 March 1994.

**II. *Oreohelix loisae*:** (1) Holotype, ANSP 407556; paratypes ANSP 408362, SBMNH 345735, Felt Wash, Goshute Mountains, Elko County, NEVADA, 40°463 N, -114°269 W, 10 May 1997. (2) Christmas Tree Canyon, Goshute Mountains, Elko County, NEVADA, 40°444 N, -114°256 W, 10 October 1986.

**III. *Oreohelix strigosa depressa*:** (1) ANSP 401886, Lutts Creek, Ruby Mountains, Elko County, NEVADA, 40°609 N, -115°319 W, 19 May 1993. (2) ANSP 401887, Horse Spring, Pilot Range, Elko County, NEVADA, 41°024 N, -114°105 W, 7 August 1992. (3) ANSP 401888, Birch Creek, Deep Creek Range, Juab County, UTAH, 14 November 1994. (4) ANSP 401890, Six Mile Creek, Pequop Range, Elko County, NEVADA, 41°003 N, -114°562 W, 4 June 1993. (5) MAP 743, Davis Canyon, Diamond Range, Eureka County, NEVADA, 40°084 N, -115°869 W, 3 November 1993. (6) MAP 868, Shingle Pass, Egan Range, Lincoln County, NEVADA, 38°522 N, -114°916 W, 25 May 1999. (7) MAP 721, Berry Creek, Schell Creek Range, White Pine County, NEVADA, 39°361 N, -114°687 W, 10 July 1993. (8) MAP 990, Snake Creek, Snake Range, White Pine County, NEVADA, 38°994 N, -114°240 W, 13 May 1989. (9) MAP 820, Hendrys Creek, North Snake Range, White Pine County, NEVADA, 39°227 N, -114°136 W, 14 May 1988. (10) MAP 832, Goshute Creek, Cherry Creek Range, White Pine County, NEVADA, 40°058 N, -115°835 W, 8 August 1988. (11) MAP 747, Ackler Creek, East Humboldt Range, Elko County, NEVADA, 40°996

N, -115°197 W, 2 June 1989. (12) MAP 818, southwest of Swasey Peak, House Range, Millard County, UTAH.

**IV. *Oreohelix hemphilli*:** (1) SBMNH 345737, Cathedral Canyon, Mount Hamilton, White Pine Range, White Pine County, NEVADA, 39°212 N, -115°465 W, 20 July 2000. (2) MAP 761, Murphy Wash, south Snake Range, White Pine County, NEVADA, 38°788 N, -114°308 W, 28 May 1997. (3) MAP 669, Pearl Creek, southern Ruby Mountains, Elko County, NEVADA, 40°271 N, -115°551 W, 11 September 1999. (4) MAP 806, Smith Creek, north Snake Range, White Pine County, NEVADA, 39°163 N, -114°164 W, 28 March 1994. (5) MAP 829, Taylor Canyon, Cherry Creek Range, Elko County, NEVADA, 40°231 N, -114°943 W, 21 July 1995. (6) MAP 700, Hooper Canyon, Quinn Canyon Range, Nye County, NEVADA, 38°236 N, -115°661 W, 8 April 1993. (7) MAP 433, Kingston Canyon, Toiyabe Mountains, Lander County, NEVADA, 39°204 N, -117°137 W, 10 July 1998. (8) MAP 713, Felt Wash, Goshute Range, Elko County, NEVADA, 40°463 N, -114°269 W, 10 May 1997. (9) MAP 872, Roberts Creek, Roberts Creek Mountains, Eureka County, NEVADA, 39°811 N, -116°312 W, 25 July 1998. (10) MAP 989, canyon on west slope of Highland Range, Lincoln County, NEVADA, 37°855 N, -114°591 W, 26 May 2000. (11) MAP 914, Pine Creek, Toiyabe Range, Nye County, NEVADA, 38°786 N, -116°883 W, 30 June 1999. (12) MAP 743, Six Mile Creek, Pequop Range, Elko County, NEVADA, 40°319 N, -114°553 W, 3 August 1993. (13) MAP 74, Hendrys Creek, north Snake Range, White Pine County, NEVADA, 39°228 N, -114°136 W, 16 May 1987. (14) MAP 905, Holt Camp Canyon, Egan Range, White Pine County, NEVADA, 39°704 N, -114°884 W, 17 September 1988. (15) MAP 693, Davis Canyon, Diamond Range, Eureka County, NEVADA, 41°084 N, -115°869 W, 3 November 1993. (16) MAP 909, Scofield Canyon, Grant Range, Nye County, NEVADA, 38°302 N, -115°508 W, 15 June 1999. (17) MAP 857, Hogan's Canyon, Pilot Range, Box Elder County, UTAH, (18) MAP 1002, Birch Creek, Deep Creek Range, Juab County, UTAH. (19) MAP 1073 Miller Canyon, House Range, Millard County, UTAH.