



2-20-2004

Groundwater invertebrate fauna of the Bear River Range, Utah

Candace Brindza Huebner
Utah State University, Logan

Mark R. Vinson
Utah State University, Logan

Follow this and additional works at: <https://scholarsarchive.byu.edu/wnan>

Recommended Citation

Huebner, Candace Brindza and Vinson, Mark R. (2004) "Groundwater invertebrate fauna of the Bear River Range, Utah," *Western North American Naturalist*. Vol. 64 : No. 1 , Article 16.
Available at: <https://scholarsarchive.byu.edu/wnan/vol64/iss1/16>

This Note is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in Western North American Naturalist by an authorized editor of BYU ScholarsArchive. For more information, please contact scholarsarchive@byu.edu, ellen_amatangelo@byu.edu.

GROUNDWATER INVERTEBRATE FAUNA OF THE BEAR RIVER RANGE, UTAH

Candace Brindza Huebner¹ and Mark R. Vinson¹

Key words: groundwater, aquatic invertebrates, *Stygobromus*, Utah.

The chance discovery of an undescribed species of blind subterranean amphipod within the *Stygobromus hubbsii* group in a cave in northern Utah prompted us to question the possible distribution and occurrence of additional groundwater fauna in this area. Groundwater biota in western North America have not been particularly well studied (cf. Stanford and Gauffin 1974, Ward 1977, Pennak and Ward 1986, Stanford and Ward 1988, Stanford et al. 1994, Ward and Voelz 1994, Ward et al. 1994, Drost et al. 1997, Canton and Chadwick 2000), especially when compared with studies of western surface water systems or groundwater systems of eastern North America, where karst topography is more prevalent overall and less disjunct (Ritter 1986, also see Culver 1997 [<http://www.utexas.edu/depts/tnhc/.www/biospeology/uskarst.jpg>]). The objective of our study was to follow up on this chance find by more intensively sampling the aquatic macroinvertebrates present in this area of northern Utah.

This study was conducted in Cache Valley and the Bear River Mountain Range in Cache County, Utah. Cache Valley is an elongated graben formed by normal high-angle faults typical of basins in the Basin and Range Province (Fenneman 1931). The Bear River Mountains, which are immediately east of Cache Valley, are composed of sedimentary and metamorphic rock of Permian to Precambrian age, including limestone, dolomite, quartzite, sandstone, mudstone, siltstone, and shale (Kariya et al. 1994). Groundwater in Cache Valley and the Bear River Mountain Range, originating from infiltration of precipitation and seepage from streams, is considered a shallow and unconfined, unglaciated karst system (Kariya et al. 1994).

Sampling locations included a stream flowing within Logan Cave and 27 spring outflows (Table 1). These springs generally flow directly or indirectly from fractures, faults, joints, and solution channels in carbonate rocks (Bjorklund and McGreevy 1971). Sixty spring sites were initially identified for sampling from topographic maps, but after visiting each spring we excluded many from the study because of lack of water or extensive human alteration, e.g., dewatering caused by construction of live-stock watering ponds. Most studies of groundwater fauna have been based on collections from either wells or caves, but spring outflows also can be used as collection sites of groundwater organisms (Nielsen 1950, Erman et al. 1995).

We sampled aquatic invertebrates at 5 locations in fall 1998, and all 28 sites were sampled at various times between May and November 1999. The stream within Logan Cave was sampled at several locations approximately 400–1000 m from the cave entrance. Springs were sampled at their point of discharge. Within Logan Cave we set 3 drift nets (10-cm diameter, 250-micron mesh) in the stream at approximately mid-depth. At each spring location, a short length (0.5 m) of 10- or 16-cm-diameter polyvinyl chloride (PVC) pipe, depending on spring outflow volume, was driven about 0.3 m into the hillside where the spring exited the ground. These pipes limited collection of downstream surface water organisms by capturing 100% of the flow at each spring. A 250-micron nylon mesh bag installed over the pipe outlet filtered all water exiting each spring. Nets were left in place at all sites for 1 week, after which all material collected in the nets was preserved in 70% ethanol and returned to the laboratory.

¹Department of Aquatic, Watershed, and Earth Resources, Utah State University, Logan, UT 84322-5210.

TABLE 1. Location of sampling sites within the Bear River Range and number of *Stygobromus* collected at each site.

Sampling location	Latitude (N)	Longitude (W)	County, State	<i>Stygobromus</i> collected
Logan Cave, 3 sampling locations	41.8000	111.6250	Cache County, Utah	28
Beaver Mountain Spring	41.9611	111.5342	Cache County, Utah	5
Beaver Spring	41.9545	111.5857	Cache County, Utah	1
Box Spring	41.7265	111.6240	Cache County, Utah	
Card Canyon Spring	41.7530	111.6520	Cache County, Utah	1
Coldwater Spring	41.7178	111.5998	Cache County, Utah	
Coldwater Spring by Tony Grove	41.8822	111.6462	Cache County, Utah	
Franklin Basin, unnamed spring # 2	41.9615	111.5952	Cache County, Utah	
Franklin Basin, unnamed spring # 3	41.9745	111.5987	Cache County, Utah	3
Grey Cliff Spring	41.6731	111.6173	Cache County, Utah	
Hodges Spring	41.9806	111.4851	Cache County, Utah	
Millville Spring	41.6587	111.7698	Cache County, Utah	
Mosslander Spring	41.6578	111.6199	Cache County, Utah	7
North Cheney Spring	41.8930	111.4683	Cache County, Utah	
Pig Hole Spring	41.6543	111.6355	Cache County, Utah	4
Pine Spring	41.7288	111.6172	Cache County, Utah	
Pleasant Val Spring	41.6477	111.5787	Cache County, Utah	
Pole Hole Spring	41.6590	111.5713	Cache County, Utah	
Porcupine Reservoir Spring	41.5190	111.7545	Cache County, Utah	2
Rock Spring	41.7863	111.5395	Cache County, Utah	
Sadducee Spring	41.0513	111.4600	Rich County, Utah	
Sow Hole Spring	41.6561	111.6068	Cache County, Utah	
Spring at Spring Campground	41.6333	111.6500	Cache County, Utah	
Spring Hollow	41.7572	111.7188	Cache County, Utah	
St. Charles Spring	42.1120	111.4473	Bear Lake County, Idaho	4
Upper Sow Hole Spring	41.6560	111.6047	Cache County, Utah	1
Wind Cave	41.7625	111.7035	Cache County, Utah	
Worm Fence Spring	41.8597	111.5500	Cache County, Utah	1

All organisms from each sample were removed and identified to the lowest possible taxonomic level. Dr. John Holsinger (Old Dominion University, VA) verified amphipod identifications.

Habitat conditions in Logan Cave and at the point where samples were collected at the spring sites were similar. Substrate consisted of gravel with some silt deposits. Organic matter was present at all sites and consisted primarily of roots, small branches, and fine debris. Organic matter within Logan Cave was visibly less than that found downstream of the collection points at the spring sites, as there were no larger pieces of wood or moss growing within the cave. Water temperatures were highly similar among all sites and ranged from 5°C to 7°C. Ranging from 7.1 to 8.4, pH was also fairly similar among sites. Specific conductance within Logan Cave was 315 $\mu\text{S cm}^{-1}$ near the mouth of the cave and 359 $\mu\text{S cm}^{-1}$ deeper into the cave. Specific conductance at spring sites was 160–525 $\mu\text{S cm}^{-1}$ and averaged 305 $\mu\text{S cm}^{-1}$.

From the 28 sites, we collected and identified 6443 individuals and 44 genera of aquatic macroinvertebrates (Table 2). Most individuals were small and immature, which made identifications beyond family and genera difficult if not impossible. Invertebrate genera richness varied from 2 to 17 at individual sites. No aquatic vertebrates were collected. We did, however, collect an undescribed species within the *Stygobromus hubbsii* group (Dr. John Holsinger personal communication) from 12 spring sites and Logan Cave. The most frequently collected organisms were Chloroperlidae, Capniidae, Collembola, Dytiscidae, *Ameletus*, and *Stygobromus hubbsii* group. All taxa collected in Logan Cave also were collected at spring outflows.

The invertebrate fauna we collected is similar to that collected in Roaring Springs Cave, Grand Canyon, Arizona, by Drost and Blinn (1997). *Stygobromus* is the only obligate groundwater fauna we collected; the other taxa are considered occasional hyporheic dwellers (Boulton 2000) and are commonly found in the Logan River (Vinson unpublished data). Since

TABLE 2. List of taxa and number of individuals collected in Logan Cave (indicated by an *) and 28 spring outflows in the Bear River Mountain Range, northern Utah.

Taxon	Individuals collected	Taxon	Individuals collected
Annelida		Ephemeroptera	
Oligochaeta	95	Ameletidae	
Arthropoda		<i>Ameletus</i> *	20
Arachnida		Baetidae	36
Trombidiformes*	673	<i>Baetis</i>	901
Entognatha		Ephemerellidae	4
Collembola*	37	Heptageniidae, unidentified	9
Diplura*	2	<i>Cinygmula</i>	5
Insecta		<i>Epeorus</i>	3
Coleoptera	3	Hemiptera	
Curculionidae	2	Gerridae	
Dytiscidae, unidentified	9	<i>Aquarius</i>	1
<i>Agabus</i>	75	Lepidoptera, unidentified	1
<i>Deronectes</i>	4	Plecoptera	
<i>Hydroporus</i>	311	Capniidae, unidentified*	35
<i>Hydrocatus</i>	29	Chloroperlidae	
<i>Oreodytes</i>	1	<i>Sweltsa</i> *	32
Elmidae		Nemouridae, unidentified	54
<i>Heterlimnius</i>	22	<i>Malenka</i>	134
<i>Narpus</i>	3	<i>Zapada</i>	260
<i>Ordobrevia</i>	1	Perlidae	
<i>Stenelmis</i>	2	<i>Hesperoperla pacifica</i>	34
Hydraenidae		Perlodidae	215
<i>Hydraena</i>	1	<i>Isoperla</i>	761
Hydrophilidae		Trichoptera	192
<i>Ametor</i>	10	Hydroptilidae, unidentified	13
<i>Paracymus</i>	7	Limnephilidae, unidentified	12
Diptera	2	<i>Dicosmoecus</i>	1
Ceratopogonidae, unidentified	5	<i>Hesperophylax</i>	480
<i>Probezzia</i>	3	<i>Psychoglypha</i>	48
Chironomidae, unidentified	1006	Rhyacophilidae	
Dixidae		<i>Rhyacophila</i>	5
<i>Dixa</i>	19	Uenoidae	
Empididae		<i>Neophylax</i>	8
<i>Clinocera</i>	4	<i>Neothremma</i>	296
Ephydriidae	1	<i>Oligophlebodes</i>	1
Muscidae	19	Malacostraca	
Psychodidae		Amphipoda	
<i>Pericoma</i>	2	Crangonyctidae	
Simuliidae		<i>Stygobromus hubbsi</i> group*	52
<i>Simulium</i>	58	Mollusca	
Stratiomyidae		Bivalvia	
<i>Caloparyphus</i>	10	Veneroidea	
<i>Euparyphus</i>	9	Pisidiidae	
Tabanidae		<i>Pisidium</i>	3
<i>Tabanus</i>	2	Gastropoda	
Tipulidae, unidentified	1	Basommatophora	
<i>Dicranota</i>	17	Lymnaeidae, unidentified	9
<i>Limonia</i>	3	Planorbidae, unidentified	8
<i>Ormosia</i>	10	Neotaenioglossa	
<i>Pedicia</i>	1	Hydrobiidae, unidentified	18
<i>Tipula</i>	2	Platyhelminthes	
		Turbellaria	331

this study was performed, we also have collected *Stygobromus* in southern Utah (Alvey Wash, Garfield County). These discoveries were predicted by Holsinger (1974) and echoed by Ward (1977) and Canton and Chadwick (2000). The occurrence and discovery of additional species of *Stygobromus* in other areas of the western United States are likely as these areas are sampled in the future.

We thank Dr. John Holsinger for examination of the *Stygobromus* specimens. Charles Hawkins offered valuable suggestions regarding the study and the manuscript. We also thank B.C. Kondratieff for reviewing the manuscript. Financial support was provided by Utah State University.

LITERATURE CITED

- BJORKLUND, L.J., AND L.J. MCGREEVY. 1971. Ground-water resources of Cache Valley, Utah and Idaho. Utah Department of Natural Resources Technical Publication 36. 72 pp.
- BOULTON, A. 2000. The subsurface macrofauna. Pages 337–361 in J. Jones and P. Mulholland, editors, Streams and groundwaters. Academic Press, San Diego, CA.
- CANTON, S.P., AND J.W. CHADWICK. 2000. Distribution of the subterranean amphipod *Stygobromus* in central Colorado streams, with notes on the interstitial community. *Western North American Naturalist* 60:130–138.
- CULVER, D. 1997. Color map of the karst regions of the USA. Presented at the Conference on the Conservation and Protection of the Biota of Karst, Nashville, TN.
- DROST, C.A., AND D.W. BLINN. 1997. Invertebrate community of Roaring Springs Cave, Grand Canyon National Park, Arizona. *Southwestern Naturalist* 42: 497–500.
- ERMAN, N.A., AND D.C. ERMAN. 1995. Spring permanence: Trichoptera species richness, and the role of drought. *Journal of the Kansas Entomological Society* 68: 50–64.
- FENNEMAN, N.M. 1931. Physiography of the western United States. McGraw-Hill, New York. 534 pp.
- HOLSINGER, J.R. 1974. Systematics of the subterranean amphipod genus *Stygobromus* (Gammaridae), part I: species of the western United States. *Smithsonian Contributions to Zoology* 160:1–63.
- KARIYA, K.A., D.M. ROARK, AND K.M. HANSON. 1994. Hydrology of Cache Valley, Cache County, Utah, and adjacent part of Idaho, with emphasis on simulation of ground-water flow. Utah Department of Natural Resources Technical Publication 108. 120 pp.
- NIELSEN, A. 1950. On the zoogeography of springs. *Hydrobiologia* 2:313–320.
- PENNAK, R.W., AND J.V. WARD. 1986. Interstitial faunal communities of the hyporheic and adjacent ground-water biotopes of a Colorado mountain stream. *Archiv für Hydrobiologie Supplement* 74:356–396.
- ITTER, D.F. 1986. Process geomorphology. 2nd edition. Wm. C. Brown Publishers, Dubuque, IA. 579 pp.
- STANFORD, J.A., AND R. GAUFIN. 1974. Hyporheic communities of two Montana rivers. *Science* 18:700–702.
- STANFORD, J.A., AND J.V. WARD. 1988. The hyporheic habitat of river ecosystems. *Nature* 335:64–66.
- STANFORD, J.A., ET AL. 1994. Alluvial aquifers of the Flathead River, Montana. Pages 367–390 in J. Gilbert, D.L. Danielopol, and J.A. Stanford, editors, Groundwater ecology. Academic Press, San Diego, CA.
- WARD, J.V. 1977. First records of subterranean amphipods from Colorado with descriptions of three new species of *Stygobromus* (Crangonyctidae). *Transactions of the American Microscopical Society* 96:452–466.
- WARD, J.V., AND N.J. VOELZ. 1994. Groundwater fauna of the South Platte River system, Colorado. Pages 391–423 in J. Gilbert, D.L. Danielopol, and J.A. Stanford, editors, Groundwater ecology. Academic Press, San Diego, CA.
- WARD, J.V., J.A. STANFORD, AND N.J. VOELZ. 1994. Spatial patterns of Crustacea in the floodplain aquifer of an alluvial river. *Hydrobiologia* 287:11–17.

Received 22 January 2002

Accepted 28 March 2003