



4-24-2009

New distribution record for the social parasitic ant *Anergates atratulus* (Schenck, 1852) (Hymenoptera: Formicidae): an ICUN red-listed species

Shawn T. Dash

University of Texas at El Paso, stdash@miners.utep.edu

Leticia Sanchez

University of Colorado at Boulder

Follow this and additional works at: <https://scholarsarchive.byu.edu/wnan>

Recommended Citation

Dash, Shawn T. and Sanchez, Leticia (2009) "New distribution record for the social parasitic ant *Anergates atratulus* (Schenck, 1852) (Hymenoptera: Formicidae): an ICUN red-listed species," *Western North American Naturalist*: Vol. 69 : No. 1 , Article 19.
Available at: <https://scholarsarchive.byu.edu/wnan/vol69/iss1/19>

This Note is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in Western North American Naturalist by an authorized editor of BYU ScholarsArchive. For more information, please contact scholarsarchive@byu.edu, ellen_amatangelo@byu.edu.

NEW DISTRIBUTION RECORD FOR THE SOCIAL PARASITIC ANT
ANERGATES ATRATULUS (SCHENCK, 1852) (HYMENOPTERA:
FORMICIDAE): AN IUCN RED-LISTED SPECIES

Shawn T. Dash¹ and Leticia Sanchez²

ABSTRACT.—The International Union for Conservation of Nature and Natural Resources red-lists a number of ant species. One species is *Anergates atratulus* (Schenck, 1852), a rarely collected social parasite introduced from Europe. The known distribution in North America is along the Atlantic seaboard. This note reports a huge range expansion in North America, to include Colorado. This finding is important because it provides useful information about the conservation and dispersal patterns of a rare, introduced, socially parasitic ant species.

Key words: black guest ant, *Anergates atratulus*, *Tetramorium caespitum*, Colorado, introduced species, dispersal.

The black guest ant *Anergates atratulus* (Schenck, 1852) is rarely encountered (Fisher and Cover 2007), perhaps because of its socially parasitic lifestyle or merely because it is indeed just rare (Wheeler 1910, Hölldobler and Wilson 1990). To date, these obligate social parasites have been found only within nests of *Tetramorium caespitum* (L., 1758) in North America (Creighton 1950, Skinner 1987, Fisher and Cover 2007). *Anergates atratulus* is a workerless parasite that is fed exclusively by *T. caespitum* workers (Wheeler 1910, Creighton 1950, Skinner 1987). The natural history of *A. atratulus* is complex and interesting and may provide insight into the distribution of the species. After adelphogametic (mating of brother and sisters) fertilization within the host nest, mated females disperse. At the completion of this dispersal flight, the female locates a nest of the host species and enters a weakened colony, unmolested by *T. caespitum* workers. (Schenck 1852, Wheeler 1910, Hölldobler and Wilson 1990). Nests of *T. caespitum* parasitized by *A. atratulus* consist of a few workers of *T. caespitum*, a number of virgin *A. atratulus* females, and pupoidal *A. atratulus* males (Wheeler 1910). One may assume that a certain density and demographic of *T. caespitum* nests are needed for *A. atratulus* to colonize.

Both *A. atratulus* and *T. caespitum* are widely distributed in Europe and are thought to have been introduced into North America

during the colonial period (Smith 1943, Brown 1957, Czechowski et al. 2002, Ward 2005). Bolton (1979) reported that *T. caespitum* ranges from the eastern United States west to Washington and California but that the western distribution is localized. To date, *A. atratulus*, in contrast, has been recorded only in the New England and mid-Atlantic states: Connecticut, Pennsylvania, New Jersey, Maryland, Delaware, Ohio, Washington DC, and Virginia (Smith 1979, Coovert 2003, Fisher and Cover 2007).

We report the first record of *Anergates atratulus* from the western United States. A single female was collected in a pitfall trap near Niwot in eastern Boulder County, on Niwot Trail close to the Boulder and Lefthand irrigation systems near the 79th Street trailhead (40°05.447 N, 105°09.887 W at 1555.85 m on 7 June 2005). The specimen collected in this study will be deposited in the holdings at the University of Colorado at Boulder (UCMC). The habitat is high-elevation grassland characterized by native and nonnative grasses, including 2 species of *Bromus* as well as Kentucky bluegrass (*Poa arachnifera*), interspersed with riparian vegetation dominated by cottonwoods and willows. The traps were 30–50 m from the riparian vegetation. *Tetramorium caespitum* workers were found in 39% of pitfall traps located about 10 m from paved trails (where nests are most abundant). A low abundance of workers was found in all traps. Gregg

¹Laboratory for Environmental Biology, Dept. of Biological Sciences, Centennial Museum, University of Texas at El Paso, El Paso TX 79968. E-mail: stdash@miners.utep.edu

²Ecology and Evolutionary Biology, University of Colorado at Boulder, Boulder, CO 80309.

(1963) did not record *T. caespitum* in Colorado. However, by the mid 1990s the species was a major component of many communities in Colorado (W. Cranshaw personal communication). We found, based on museum records (CSUMC), that the first specimens from Colorado were collected in 1985. *Tetramorium caespitum* is now the major ant pest in homes in Colorado and has become a component of ant communities.

The range expansion reported here represents the westernmost record for *A. atratulus*. Given the obligate relationship between *A. atratulus* and *T. caespitum*, *A. atratulus* was not likely to have successfully invaded the area until *T. caespitum* was well established. Because *A. atratulus* is listed by the IUCN as a vulnerable taxon (Social Insects Specialist Group 1996), this range expansion has important conservation implications. Most notably, it could be that the range of *A. atratulus* is much larger than previously suspected but that where *A. atratulus* occurs, it is rare. However, in the case of *A. atratulus*, it is more likely that rarity is a result of limited sampling. Generally speaking, populations of social parasites are smaller and the ranges more restricted than those of their host species. This is likely the case with *A. atratulus* and *T. caespitum*.

Appreciation is given to William Mackay, Israel Del Toro, and Michael Breed for reviewing and providing comments on early versions of this note. We also thank Nathan Sanders and an anonymous reviewer for providing comments and critiques. Funding was provided by University of Colorado Museum of Natural History's Walker van Riper Grant. This material is based upon work supported by the National Science Foundation under Grant No. 0405470, W. Mackay, principal investigator. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

LITERATURE CITED

- BOLTON, B. 1979. The ant tribe Tetramoriini (Hymenoptera: Formicidae), the genus *Tetramorium* Mayr in the Malagasy region and in the New World. *Bulletin of the British Museum of Natural History (Entomology)* 38:129–181.
- BROWN, W.L. 1957. Is the ant genus *Tetramorium* native in North America? *Breviora* 72:1–8.
- CREIGHTON, W.S. 1950. *Ants of North America*. Bulletin of the Museum of Comparative Zoology 104. 585 pp., 57 plates.
- COOVERT, G.A. 2003. The ants of Ohio (Hymenoptera: Formicidae). *Bulletin of the Ohio Biological Survey* 15. 196 pp.
- CZECHOWSKI, W., A. RADCHENKO, AND W. CZECHOWSKA. 2002. The ants (Hymenoptera: Formicidae) of Poland. Museum and Institute of Zoology, Warszawa, Poland. 200 pp.
- FISHER, B.L., AND S.P. COVER. 2007. *Ants of North America, a guide to genera*. University of California Press, Berkeley. 194 pp.
- GREGG, R.E. 1963. The ants of Colorado, with reference to their ecology, taxonomy, and geographic distribution. University of Colorado Press, Boulder. 792 pp.
- HÖLDOBLER, B., AND E.O. WILSON. 1990. *The ants*. Harvard University Press, Cambridge, MA. 732 pp.
- SCHENCK, C.F. 1852. Beschreibung Nassauischer Ameisenarten. *Jahrbuch. Der Vereinigung für Naturkunde. Herzogthum Nassau Wiesbaden* 8:1–149.
- SKINNER, G.J. 1987. *Ants of the British Isles*. Shire Natural History, Number 21. Shire Publications Ltd., Aylesbury, U.K.
- SMITH, D.R. 1979. Family Formicidae. Pages 1323–1467 in K.V. Krombein, P.D. Hurd, Jr., D.R. Smith, and B.D. Burks, editors, *Catalog of Hymenoptera in America North of Mexico*. Volume 2. Smithsonian Institution Press, Washington, DC. 1010 pp.
- SMITH, M.R. 1943. Ants of the genus *Tetramorium* in the United States with the description of a new species. *Entomological Society of Washington* 45:1–5.
- SOCIAL INSECTS SPECIALIST GROUP. 1996. *Anergates atratulus*. In: 2006 IUCN Red List of Threatened Species. Available from: <http://www.iucnredlist.org>
- WARD, P.S. 2005. Synoptic review of the ants of California (Hymenoptera: Formicidae). *Zootaxa* 936:1–68.
- WHEELER, W.M. 1910. *Ants, their structure, development, and behavior*. Columbia University Press, New York. 663 pp.

Received 26 September 2007

Accepted 12 September 2008