

## RANGE EXTENSION OF THE GIANT WATER BUG *BELOSTOMA FLUMINEUM* SAY 1832 (HEMIPTERA: BELOSTOMATIDAE) TO SASKATCHEWAN, CANADA

Iain D. Phillips<sup>1,2,3,6</sup>, Stephen Srayko<sup>2,3</sup>, Kate S. Prestie<sup>2,3,5</sup>, Aaron J. Bell<sup>3</sup>, and Dale Parker<sup>4</sup>

**ABSTRACT.**—We collected a single specimen of the giant water bug *Belostoma flumineum* Say 1832 (Hemiptera: Belostomatidae) during routine macroinvertebrate biomonitoring in the Souris River, Saskatchewan. This is the first record of *B. flumineum* in the province of Saskatchewan and represents a more northern record than the previously documented locations in Montana, North Dakota, and southeastern Manitoba.

**RESUMEN.**—Colectamos un único ejemplar de la chinche gigante de agua *Belostoma flumineum* Say 1832 (Hemiptera: Belostomatidae) durante un biomonitoreo de rutina de macroinvertebrados en el río Souris, Saskatchewan. Este es el primer registro de *B. flumineum* en la provincia de Saskatchewan y representa un récord del norte en lugares previamente documentados de Montana, Dakota del Norte y el sureste de Manitoba.

Monitoring programs that study the impacts of anthropogenic inputs and stress on aquatic systems form the basis of most aquatic macroinvertebrate research in Saskatchewan (MoE and SWA 2012). These programs traditionally focus on single-season sampling regimes and are thus limited in the ability to detect some species with distinct seasonality in their life stages (e.g., summer diapause in Hoemsen et al. 2015). Nevertheless, recent monitoring has produced a number of new species records for the province of Saskatchewan (Parker and Phillips 2007, Phillips et al. 2008, 2013, Hoemsen et al. 2015), highlighting understudied regions of biodiversity and strengthening the understanding of biogeographical links between regions (Lehmkuhl 1980, Hoemsen et al. 2015). This new information suggests that the single-season sampling approach to macroinvertebrate sampling may underestimate the overall diversity of aquatic invertebrates in the province.

An adult specimen of the giant water bug *Belostoma flumineum* Say 1832 (Hemiptera: Belostomatidae) was collected on 10 September 2015 in the Souris River (49°4'46.75"N, 102°45'9.69"W; Fig. 1) near the town of Roche Percée, Saskatchewan, Canada. The specimen was collected using a 500- $\mu$ m-mesh D-frame net as part of routine biomonitoring of the

Souris River watershed by the Benthic Entomology Lab of the Water Security Agency. The voucher specimen is retained in the collection at the Water Security Agency of Saskatchewan. This specimen represents the first record of this species in the province of Saskatchewan, as well as an extension of the species' previously known distribution (see Fig. 1) in the Northern Great Plains. No previous records of *B. flumineum* exist from Saskatchewan as determined by consultation with the Royal Saskatchewan Museum (Cory Sheffield, personal communication), Agriculture and Agri-Food Canada (Tyler Wist, personal communication), and Aquatax Consulting (Dale Parker, personal communication).

The Souris River in this region passes through a scrubland and forested valley within aspen parkland, and the reach is characterized by sediments of clay and silt with an abundance of woody debris and macrophytes. Abiotic characteristics of the site at the time of collection featured a velocity of approximately  $0.13 \text{ m} \cdot \text{s}^{-1}$ , specific conductivity of  $1715 \text{ } \mu\text{S} \cdot \text{cm}^{-1}$ , pH of 7.8, turbidity of 127.5 NTU, and dissolved oxygen concentration of  $6.35 \text{ mg} \cdot \text{L}^{-1}$ .

*Belostoma* is one of 4 genera of Belostomatidae recorded in North America. Nine species of *Belostoma* occur in North America, including

<sup>1</sup>Water Quality and Habitat Assessment Services, Water Security Agency, #101–108 Research Drive, Saskatoon, Saskatchewan, Canada S7N 3R3.

<sup>2</sup>Department of Biology, University of Saskatchewan, #112 Science Place, Saskatoon, Saskatchewan, Canada S7N 5E2.

<sup>3</sup>Troutreach Saskatchewan, Saskatchewan Wildlife Federation, #9 Lancaster Road, Moose Jaw, Saskatchewan, Canada S7J 1M8.

<sup>4</sup>AquaTax Consulting, #1204 Main Street, Saskatoon, Saskatchewan, Canada S7H 0L2.

<sup>5</sup>Toxicology Centre, University of Saskatchewan, 44 Campus Drive, Saskatoon, Saskatchewan, Canada S7N 5B3.

<sup>6</sup>Corresponding author. E-mail: iain.phillips@wsask.ca




Fig. 1. Known distribution of *Belostoma flumineum* showing the new record of this species in Saskatchewan (green star) and previously documented collection locations in Montana, North Dakota, and Manitoba (black circles). Green circles indicate areas that were sampled as part of a larger biomonitoring program in 2006–2013 during which this species was not detected.



Fig. 2. Belostomatid taxa that occur in Saskatchewan: *Lethocerus americanus* Leidy 1847, 50 mm (above); and the new provincial record, *Belostoma flumineum*, 21 mm (below). Scale bar: 10 mm.

*B. bakeri* Montandon, *B. confusum* Lauck, *B. ellipticum* Latreille, *B. flumineum*, *B. fusciven-*

*tre* (Dufour), *B. lutarium* (Stål), *B. saratogae* Menke, *B. subspinosum* (Palisot), and *B. testaceum* (Leidy) (Menke 1979). In the United States, *B. flumineum* occurs from New England to Florida, as well as in Louisiana, Colorado, and California (Brooks and Kelton 1967, Menke 1979). *Belostoma flumineum* has been recorded throughout Canada, from Newfoundland to Manitoba in the east (Brooks and Kelton 1967), and British Columbia in the west (Downes 1934). However, despite significant search efforts, *B. flumineum* has not been documented in the Northern Great Plains west of Manitoba (Brooks and Kelton 1967, Parker and Phillips 2007). In a recent search, Scudder et al. (2010) conducted extensive sampling of Hemiptera in 50 water bodies across the Prairie Provinces, including a variety of lakes and ponds, and found *B. flumineum* in only 5 forested areas in southeastern Manitoba. Similarly, previous monitoring efforts in southeastern Saskatchewan and around the Souris River area (Fig. 1) from 2006 to 2013 failed to detect this species.

The feeding and reproductive habits of belostomatids are well known and summarized in Smith (1975), Menke (1979), and Polhemus (1984). In short, belostomatids occupy shallow areas in bodies of standing water, as well as pools and marginal areas in lotic habitats (Merritt et al. 2008). Due to their relatively large size (Fig. 2), belostomatids can act as top predators in the absence of vertebrates, thereby influencing ecosystem function and shaping trophic structure and energy flow within aquatic food webs (Cooper 1983, Runck and Blinn 1994).

#### LITERATURE CITED

- BROOKS, A.R., AND L.A. KELTON. 1967. Aquatic and semiaquatic Heteroptera of Alberta, Saskatchewan, and Manitoba (Hemiptera). *Memoirs of the Entomological Society of Canada* 51:1–91.
- COOPER, S.D. 1983. Selective predation on cladocerans by common pond insects. *Canadian Journal of Zoology* 61:879–886.
- DOWNES, W. 1934. Additions to the list of B. C. Hemiptera. *Proceedings of the Entomological Society of British Columbia* 31:46–48.
- HOEMSEN, B.M., I.D. PHILLIPS, D.W. PARKER, A.J. BELL, J.A. BERGSVEINSON, J.S. ARMSTRONG, AND D.P. CHIVERS. 2015. Extended family: a caddisfly new to Saskatchewan, Canada with notes on the life history of *Neophylax splendens* (Trichoptera: Thremmatidae). *Canadian Entomologist* 147:425–431.
- LEHMKUHL, D.M. 1980. Temporal and spatial changes in the Canadian insect fauna: patterns and explanation in the prairies. *Canadian Entomologist* 112:1145–1159.
- MENKE, A.E., EDITOR. 1979. The semiaquatic and aquatic Hemiptera of California (Heteroptera: Hemiptera). *Bulletin of the California Insect Survey* 21:1–666.
- MERRITT, R.W., K. CUMMINS, AND M.B. BERG. 2008. *Aquatic insects of North America*. 4th edition. Kendall/Hunt Publishing Company, Dubuque, IA.
- [MOE AND SWA] MINISTRY OF ENVIRONMENT AND SASKATCHEWAN WATERSHED AUTHORITY. 2012. *Saskatchewan Northern Great Plains Ecosystem Health Assessment Manual 2012*. Saskatchewan Watershed Authority, Saskatoon, Saskatchewan, Canada.
- PARKER, D., AND I. PHILLIPS. 2007. Collection records of three aquatic bugs (Heteroptera); pygmy backswimmer (Pleidae), water scorpion (Nepidae) and marsh treader (Hydrometridae) for Saskatchewan, Canada. *Blue Jay* 65:143–148.
- PHILLIPS, I.D., D. PARKER, B.M. HOEMSEN, A.J. BELL, AND D.P. CHIVERS. 2013. Biological notes and range expansion of the non-biting midge *Odontomesa fulva* (Kieffer) (Diptera: Chironomidae). *Western North American Naturalist* 73:244–247.
- PHILLIPS, I.D., D. PARKER, AND G. MCMASTER. 2008. The aquatic invertebrate fauna of a northern prairie river: range extensions and water quality characteristics. *Western North American Naturalist* 68:173–185.
- POLHEMUS, J.T. 1984. Aquatic and semiaquatic Hemiptera. Pages 231–260 in R.W. Merritt and K.W. Cummins, editor, *An introduction to the aquatic insects of North America*. 2nd edition. Kendall/Hunt.
- RUNCK, C., AND D.W. BLINN. 1994. Role of *Belostoma bakeri* (Heteroptera) in the trophic ecology of a fishless desert spring. *Limnology and Oceanography* 39:1800–1812.
- SCUDDER, G.G.E., M.A. ALPERYN, AND R.E. ROUGHLEY. 2010. Aquatic Hemiptera of the prairie grasslands and parkland. Pages 303–323 in J.D. Shorthouse and K.D. Floate, editors, *Arthropods of Canadian grasslands: ecology and interactions in grassland habitats*. Biological Survey of Canada.
- SMITH, R.L. 1975. Bionomics and behavior of *Abedus herberti* with comparative observations on *Belostoma flumineum* and *Lethocerus medius* (Hemiptera: Belostomatidae). Doctoral dissertation, Arizona State University, Phoenix, AZ.

Received 29 November 2016  
 Accepted 4 April 2017  
 Published online 11 July 2017