



12-31-1981

Association of *Rhabdocline* needle blight and epicormic branching in Douglas-fir

Ronald M. Lanner
Utah State University

James A. Bryan
Utah State University

Follow this and additional works at: <https://scholarsarchive.byu.edu/gbn>

Recommended Citation

Lanner, Ronald M. and Bryan, James A. (1981) "Association of *Rhabdocline* needle blight and epicormic branching in Douglas-fir," *Great Basin Naturalist*: Vol. 41 : No. 4 , Article 16.
Available at: <https://scholarsarchive.byu.edu/gbn/vol41/iss4/16>

This Article is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in *Great Basin Naturalist* by an authorized editor of BYU ScholarsArchive. For more information, please contact scholarsarchive@byu.edu, ellen_amatangelo@byu.edu.

ASSOCIATION OF RHABDOCLINE NEEDLE BLIGHT AND EPICORMIC BRANCHING IN DOUGLAS-FIR

Ronald M. Lanner¹ and James A. Bryan¹

ABSTRACT.— In northern Utah, Douglas-firs (*Pseudotsuga menziesii* [Mirb.] Franco) with symptoms of *Rhabdocline* needle blight had a significantly higher frequency of epicormic branching than did healthy trees. It is not known whether *Rhabdocline* infection stimulates epicormy, or whether the proliferation of epicormics increases resistance to the disease.

Rhabdocline needle blight is a leaf disease of Douglas-fir (*Pseudotsuga menziesii* [Mirb.] Franco) caused by the ascomycete *Rhabdocline pseudotsugae* Syd. (Hepting 1971). Infection sometimes results in defoliation serious enough to cause death, even in relatively vigorous trees up to 100 years old (Davidson and Prentice 1967).

Douglas-fir, especially in the Rocky Mountains, is subject to the formation of epicormic branches that arise from dormant buds borne at the base of primary branches. The occurrence of epicormics appears to be a normal consequence of branch decline not requiring pest attack for its expression (Bryan and Lanner 1981), but the added stress of *Rhabdocline* defoliation may increase the frequency of epicormics in infected trees. The purpose of the observations reported here was to determine whether trees infected by *Rhabdocline* differ from healthy trees in their frequency of epicormic branches.

METHODS

These observations accompanied a detailed morphological study of epicormics that has been reported elsewhere (Bryan and Lanner 1981). Pole-sized Douglas-firs were selected at intervals along trails in Logan Canyon, Cache National Forest, Utah. *Rhabdocline*-infected trees were identified by the sparseness of their crowns, which had lost many of the needles predating those of the current year (N. Van Alfen, pers. comm.). These trees were found mainly in dense, pure stands on north-facing slopes between 1524 and 1860

m elevation, and often near streams. Uninfected trees in those stands were chosen randomly for comparison. Trees were examined in the summer, when needles infected in the previous year had already fallen (Davidson and Prentice 1967). Epicormic frequency was determined by examining the 10 uppermost dead primary branches below the base of the live crowns, and recording the number of those branches with associated epicormics. The Z-test (Dixon and Massey 1969) was used to test the hypothesis that the mean number of epicormics was equal for healthy and diseased trees. The hypothesis that the frequency distributions were equal was tested with the Kolmogorov-Smirnov two-sample test (Conover 1971).

RESULTS

Trees with *Rhabdocline* symptoms had an average of 4.8 (of 10) branches with epicormics in the lower crown, but similar healthy trees averaged only 3.0 such branches (Table 1). The Z-value of 2.23 was significant at the 99 percent level. The frequency distribution of trees with n number of epicormics also differed significantly ($\alpha = .05$) between healthy and diseased trees (Table 2).

DISCUSSION

It is apparent that *Rhabdocline*-infected trees exhibit a higher frequency of epicormics than do healthy trees. Our data do not indicate whether this is because repeated or periodic defoliations stimulate epicormic

¹Department of Forestry and Outdoor Recreation, Utah State University, Logan, Utah 84322. This study was funded under the McIntire-Stennis program and is published by permission of the Utah Agricultural Experiment Station as journal paper 2688.

formation, or whether trees that are *Rhabdo-
cline*-susceptible also happen to be prone to
epicormics. A third possibility is that suscep-
tible trees are more likely to survive *Rhabdo-
cline* if they balance their losses of photo-
synthetic surface area with new foliage borne
on epicormic branches. If this is the case,
*Rhabdo-
cline*-infected trees incapable of pro-
ducing numerous epicormics may have
evaded our sample by suffering higher morta-
lity. Douglas-firs in Logan Canyon, as else-
where, show marked individual variability in
their production of epicormics (Bryan and
Lanner 1981).

Johnson and Denton (1975) have credited
epicormics with promoting the survival of
Douglas-firs partially defoliated by the
spruce budworm. The survival of trees in-
fected by *Rhabdo-
cline* might also be pro-
moted by epicormics. *Rhabdo-
cline*-infected
trees may repeatedly cast off all their one-
year-old foliage the year following infection,
thus suffering a heavy loss of photosynthetic
surface. Epicormics stimulated by the in-
creased solar radiation permitted by such de-
foliation may serve as an important source of
needed photosynthate. If the sprouting of
epicormics indeed enables Douglas-fir to re-
cover from *Rhabdo-
cline* and extend its repro-
ductive life, we would expect natural selec-
tion to result in a higher frequency of that
trait where *Rhabdo-
cline* is of high frequency. According to Hepting (1971), Rocky Moun-
tain Douglas-fir (var. *glauca*) is extremely sus-
ceptible to *Rhabdo-
cline* but the Pacific coast
variety (var. *menziesii*) is relatively resistant.
Our studies (Bryan and Lanner 1981) show
that in the eastern part of the intermountain
area nearly all the Douglas-firs produce epi-
cormics, but in the western part of this re-
gion epicormic production is extremely vari-
able and is even exceeded by epicormic

frequency in coast Douglas-fir. Thus, central
Idaho plots had considerably fewer trees with
epicormics than the average of several Cali-
fornia plots, and the range was also far lower.
Therefore, unless it is demonstrated that
*Rhabdo-
cline* is a more serious pest of
Douglas-fir in the eastern intermountain re-
gion than in Idaho, our results do not support
the hypothesis that *Rhabdo-
cline* has been a
significant factor, at least at the regional lev-
el, in selection for epicormic branching. Fur-
ther study of the role of epicormics in
Douglas-fir survival and longevity seems war-
ranted, whether or not it bears a relationship
to *Rhabdo-
cline* resistance.

LITERATURE CITED

BRYAN, J. A., AND R. M. LANNER. 1981. Epicormic
branching in Rocky Mountain Douglas-fir. *Can.
J. For. Res.* 11:190-199.
CONOVER, W. J. 1971. Practical nonparametric statistics.
John Wiley and Sons, New York. 442 pp.
DAVIDSON, A. G., AND R. M. PRENTICE, eds. 1967. Impor-
tant forest insects and diseases of mutual concern
to Canada, the United States and Mexico. Dept.
For. and Rural Devel. Canada Pub. No. 1180, 248
pp.
DIXON, W. J., AND F. J. MASSEY. 1969. Introduction to
statistical analysis. McGraw-Hill Book Co., New
York.
HEPTING, G. H. 1971. Diseases of forest and shade trees
of the United States. USDA Forest Service Agri-
cultural Handbook No. 396. 658 pp.
JOHNSON, P. C., AND R. E. DENTON. 1975. Outbreaks of
the western spruce budworm in the American
northern Rocky Mountain area from 1922
through 1971. USDA Forest Service Gen. Tech.
Rep. INT-20.

TABLE 2. Frequency distribution of epicormic
branches on healthy and *Rhabdo-
cline*-infected trees.¹

Number of lateral branches with associated epicormics	Healthy trees	Diseased trees
	Percent of total	
0	15.1	4.3
1	17.8	17.4
2	15.0	8.7
3	16.5	8.7
4	12.3	8.7
5	12.3	8.7
6	1.4	13.1
7	0	17.4
8	2.7	13.0
9	5.4	0
10	1.4	0

¹Based on observations of 10 branches per tree on 73 healthy trees and 20
*Rhabdo-
cline*-infected trees.

TABLE 1. Number of branches with epicormics on
Douglas-fir trees in Logan Canyon, Utah, in relation to
*Rhabdo-
cline* infection.¹

<i>Rhabdo- cline</i> condition	Number of branches of 10		
	Mean	Range	S.D.
Healthy trees (n = 73)	3.0a	0-10	2.5
Infected trees (n = 20)	4.8a	0-8	2.9

¹Values followed by the same letter differ significantly at 0.01 level.