



6-30-1981

A checklist of the alpine vascular flora of the Teton Range, Wyoming, with notes on biology and habitat preferences

John R. Spence
Utah State University

Richard J. Shaw
Utah State University

Follow this and additional works at: <https://scholarsarchive.byu.edu/gbn>

Recommended Citation

Spence, John R. and Shaw, Richard J. (1981) "A checklist of the alpine vascular flora of the Teton Range, Wyoming, with notes on biology and habitat preferences," *Great Basin Naturalist*: Vol. 41 : No. 2 , Article 11.

Available at: <https://scholarsarchive.byu.edu/gbn/vol41/iss2/11>

This Article is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in Great Basin Naturalist by an authorized editor of BYU ScholarsArchive. For more information, please contact scholarsarchive@byu.edu, ellen_amatangelo@byu.edu.

A CHECKLIST OF THE ALPINE VASCULAR FLORA OF THE TETON RANGE, WYOMING, WITH NOTES ON BIOLOGY AND HABITAT PREFERENCES

John R. Spence^{1,2} and Richard J. Shaw¹

ABSTRACT.— A checklist of the vascular flora of the alpine zone (treeless vegetation above 9500 feet or 2900 m) of the Teton Range is presented. For each of the 216 species, six attributes are listed: flower color and shape, pollination mode, life form, habitat preference, and whether each species is found in the Arctic. White and yellow flowered species are most common, and zoophilous species greatly predominate over anemophilous and apomictic species. Perennial/biennial herbs are the most common life form. Common habitats in the alpine zone include dry and wet meadows, bogs, debris accumulations, and cliffs and rock faces. Arctic species account for 25.9 percent of the flora. The 216 species are distributed among 111 genera and 36 families. The largest families, in order of size, are Asteraceae, Poaceae, Cyperaceae, Brassicaceae, Rosaceae, and Scrophulariaceae.

The Teton Range, located for the most part inside Grand Teton National Park, is a typical fault block range. The mountains are about 60 km long and average 6 to 9 km wide. The major peaks in the center of the range and to the north are composed of Precambrian gneisses, schists, and granites. The southern peaks are capped by Paleozoic sedimentary rocks, and the divide to the west of the main peaks consists of Paleozoic and Mesozoic rocks. The range has been extensively glaciated in the past, and several small cirque glaciers of the Neoglacial age exist in sheltered areas of the range (Love and Reed 1968, Reed and Zartman 1973). The climate on the floor of Jackson Hole to the east of the Tetons is continental, with long, cold winters and a short growing season. Annual mean temperature is low. Precipitation is also low, falling mostly as snow in the winter months (Reed 1952, Shaw 1958).

The vascular flora of the Tetons has been well documented over the last 30 years (Shaw 1958, 1968, 1976). Approximately 150 species were listed as occurring in the alpine zone (Shaw 1976, unpublished data). Recent intensive collecting in portions of the alpine zone (Spence 1980) and reexamination of herbarium collections (Hartman and Lichvar 1979) have added more than 50 additional species.

The purposes of this checklist are to include all the recent information on the alpine

flora of the Teton Range and provide general information on floral characteristics, pollination modes, and habitats in the alpine zone. Phytogeographical relationships will also be mentioned. In this paper, the alpine zone is defined as high elevation areas of treeless vegetation, with the lower limits arbitrarily set at 9500 feet (2900 m). Although timberline (composed of *Pinus albicaulis*, *Picea engelmannii*, and *Abies lasiocarpa*) is generally found at elevations of 10,000 to 10,500 feet (3050 to 3200 m) in the Tetons, many areas below this support typical alpine species. Such areas include many of the cirques in the range, and the neoglacial deposits below the present glaciers.

METHODS

The checklist was collected from studies by Merkle (1962, 1963), Scott (1966), Shaw (1976), Spence (1980), and Hartman and Lichvar (1979), supplemented by personal observations and herbarium materials. All species on the list can be found on deposit at the Intermountain Herbarium (UTC) at Utah State University, the Rocky Mountain Herbarium (RM) at the University of Wyoming, and the Moose Herbarium in Grand Teton National Park. Polunin (1959) and Hultén (1968) were consulted for those species found in the Arctic. For each species six characteristics are noted. An asterisk (*) before the name of the species indicates it is found in

¹Department of Biology, UMC 45, Utah State University, Logan, Utah 84322.

²Present address: Department of Botany, University of British Columbia, Vancouver, B.C., Canada V6T 1W5.

the Arctic; a dagger (†) indicates the species is characteristically subalpine. Flower color is listed as w = white, y = yellow, v = violet, p = pink, b = blue, r = red, br = brown, g = green, and o = orange. Only species with conspicuously colored flowers are indicated. Flower shape is listed as a = actinomorphic, z = zygomorphic, and is listed only for those species with conspicuously colored flowers. Pollination mode is listed as Z = zoophilous, A = anemophilous or apomictic. Fryxell (1957), Pojar (1974), Ostler and Harper (1978), and Swales (1979) were consulted for aid in determining pollination mode. Life form is listed as s = shrub, p = perennial/biennial herb, g = perennial/biennial graminoid, and a / annual herb or graminoid. Major habitat types are listed as 1 = dry meadows, 2 = wet meadows, 3 = bogs, 4 = debris accumulations, 5 = cliffs and rock faces, 6 = neoglacial deposits. For more information on the habitat types see the discussion and Table 3.

RESULTS AND DISCUSSION

Table 1 lists the 216 species of vascular plants and presents for each species the biological attributes and habitat preferences by family in alphabetical order within the major categories Ferns, Fern Allies, and Gymnosperms; Dicots; and Monocots. The species are arranged in alphabetical order within each family. Table 2 is a statistical summary of the species and their attributes.

More than 50 percent of the species belong to the six largest families. In addition, 78 of the species belong to the 11 largest genera. It is interesting to note that the six families listed in Table 2 are also the six largest families in the flora of Teton County (Shaw 1976). Indeed, the genera *Carex*, *Potentilla*, *Poa*, *Salix*, and *Erigeron* are also listed as being among the largest genera in Teton County.

White and yellow flowered species predominate (64.1 percent). This result is similar to data presented by Ostler and Harper (1978) in a study of plant communities in the Wasatch Mountains of Utah and Idaho. In that study, from 65 to 75 percent of all conspicuously colored flowers in several alpine communities were white or yellow. This is a common feature of alpine floras throughout

the world (Wardle 1978). One possible reason for the commonness of white and yellow flowers in the alpine zone could be that the most common and important pollinators often are generalist fly and bee species, which frequently prefer such colors (Percival 1965, Moldenke 1976, Wardle 1978).

Some difficulty was encountered with the category Pollination Mode. Table 2 shows that 75.2 percent of the species are characteristically animal pollinated. This should be considered as a maximum value, because many of the species included in this category are probably autogamous, especially in the Brassicaceae. Not enough information on breeding systems of alpine species is available yet to state definitely whether a species outcrosses or is predominantly autogamous. The category Anemophilous/Apomictic species includes most monocots, *Artemisia*, *Oxyria digyna*, and the known apomicts *Taraxacum officinale* and *Polygonum viviparum* (Fryxell 1957, Swales 1979). Some of the grasses, particularly the *Poa* species, are probably partially or wholly apomictic also.

In the category Arctic and Alpine species 56, or 25.9 percent, are included. This can be compared with the Beartooth Range 150 km to the north. There, 47 percent of the species occur in the arctic as well (Johnson and Billings 1962). The lower value for the Tetons is possibly due to the somewhat drier conditions found there compared with typical Rocky Mountain ranges like the Beartooths or the Wind River Range to the southeast (Mahaney 1980). The value of 25.9 percent is intermediate between the Beartooths and northern Great Basin Ranges to the southwest of the Tetons (Billings 1978). The somewhat drier conditions, perhaps accentuated during the Hypsithermal, probably explain the absence of such widespread arctic species as *Koenigia islandica*, *Gentiana algida*, and *Saxifraga caespitosa* in the Tetons. All three species occur in the Beartooth and Wind River Ranges (Johnson and Billings 1962, Scott 1966).

Certain species, such as *Senecio integerrimus*, *Carex douglasii*, *Mitella pentandra*, and *Eriophyllum lanatum*, are more characteristic of the subalpine zone in the Tetons. The inclusion of the 18 subalpine

TABLE 1. A list of all species occurring in the alpine zone of the Teton Range with flower color and shape, pollination mode, life form, habitat preference, and origin noted. ° indicates an arctic species; † indicates a subalpine species. The attributes are listed as 1=flower color (y=yellow, w=white, v=violet, p=pink, b=blue, r=red, br=brown, g=green, and o=orange), 2=flower shape (a=actinomorphic, z=zygomorphic), 3=pollination mode (A=anemophilous or apomictic, Z=zoophilous), 4=life form (p=perennial/biennial herb, s=shrub, g=perennial/biennial graminoid, a=annual herb/graminoid), and 5=habitat (1=dry meadows, 2=wet meadows, 3=bogs, 4=debris accumulations, 5=cliffs and rock faces, 6=neoglacal deposits). For more detail see Methods.

Family	Attribute				
	1	2	3	4	5
<i>Ferns, fern allies, and gymnosperms</i>					
CUPRESSACEAE					
° <i>Juniperus communis</i> L. var. <i>depressa</i> Pursh	-	-	A	s	6
LYCOPODIACEAE					
<i>Lycopodium selago</i> L.	-	-	-	p	5
POLYPODIACEAE					
° <i>Asplenium viride</i> Huds.	-	-	-	p	4
<i>Athyrium filix-femina</i> (L.) Roth	-	-	-	p	6
<i>Cryptogramma crista</i> (L.) R.Br. ex Hook. var. <i>acrostichoides</i> (R.Br.) Clarke	-	-	-	p	4,6
° <i>Cystopteris fragilis</i> (L.) Bernh.	-	-	-	p	4,6
SELAGINELLACEAE					
<i>Selaginella densa</i> Rydb.	-	-	-	p	4,5,6
<i>Dicots</i>					
APIACEAE					
° <i>Bupleurum americanum</i> Coult. & Rose	y	a	Z	p	1,4
<i>Cymopterus hendersonii</i> (Coult. & Rose) Cronq.	y	a	Z	p	4,6
ASTERACEAE					
° <i>Achillea millefolium</i> L. ssp. <i>lanulosa</i> (Nutt.) Piper var. <i>alpicola</i> (Rydb.) Garrett	w	a	Z	p	1,4,6
† <i>Agoseris aurantiaca</i> (Hook.) Greene	o	a	Z	p	1,2
<i>A. glauca</i> (Pursh) Raf.	y	a	Z	p	1,6
† <i>Anaphalis margaritacea</i> (L.) Benth. & Hook.	w	a	Z	p	6
<i>Antennaria alpina</i> (L.) Gaertn. var. <i>media</i> (Greene) Jeps.	w	a	Z	p	1
<i>A. microphylla</i> Rydb.	w	a	Z	p	6
<i>A. umbrinella</i> Rydb.	w	a	Z	p	1,6
<i>Arnica latifolia</i> Bong.	y	a	Z	p	1,4,6
<i>A. longifolia</i> D.C. Eat.	y	a	Z	p	2,6
† <i>A. mollis</i> Hook.	y	a	Z	p	2
<i>Artemisia campestris</i> L. ssp. <i>borealis</i> (Pall.) Hall & Clements	-	-	A	s	1
° <i>A. frigida</i> Willd.	-	-	A	s	1
° <i>A. norvegica</i> Fries ssp. <i>saxatilis</i> (Bess.) Hall & Clements	-	-	A	s	4,6
<i>A. scopulorum</i> Gray	-	-	A	s	4
<i>Aster alpinus</i> (T. & G.) A. Gray var. <i>haydenii</i> (Porter) Cronq.	v	a	Z	p	1
<i>Chaenactis alpina</i> (Gray) M. E. Jones	p	a	Z	p	4,6
<i>Cirsium tweedyi</i> (Rydb.) Petr.	p	a	Z	p	6
° <i>Erigeron compositus</i> Pursh	v	a	Z	p	1,4,6
<i>E. leiomerus</i> Gray	v	a	Z	p	1,4,6
<i>E. peregrinus</i> (Pursh) Greene ssp. <i>callianthemus</i> (Greene) Cronq.	p	a	Z	p	2
<i>E. simplex</i> Greene	v	a	Z	p	1,4
<i>E. ursinus</i> D.C. Eat.	v	a	Z	p	1
† <i>Eriophyllum lanatum</i> (Pursh) Forbes var. <i>integrifolium</i> (Hook.) Smiley	y	a	Z	p	6
<i>Haplopappus acaulis</i> (Nutt.) Gray	y	a	Z	p	4,6
<i>H. suffruticosa</i> (Nutt.) Gray	y	a	Z	s	1

Table 1 continued.

Family	Attribute				
	1	2	3	4	5
<i>Hieracium gracile</i> Hook.	y	a	Z	p	2,6
<i>Hymenoxys grandiflora</i> (T. & G.) Parker	y	a	Z	p	1,4,6
<i>Senecio amplexans</i> Gray	y	a	Z	p	1,4
<i>S. canus</i> Hook.	y	a	Z	p	4
<i>S. crassulus</i> Gray	y	a	Z	p	2
<i>S. fremontii</i> T. & G.	y	a	Z	p	4,6
† <i>S. integerrimus</i> Nutt.	y	a	Z	p	6
† <i>S. streptanthifolius</i> Greene	y	a	Z	p	6
† <i>S. triangularis</i> Hook.	y	a	Z	p	2
<i>S. werneriaefolius</i> Gray	y	a	Z	p	6
* <i>Solidago multiradiata</i> Ait.	y	a	Z	p	1,4,6
* <i>Taraxacum lyratum</i> (Ledeb.) DC.	y	a	Z	p	1,4,6
<i>T. officinale</i> Weber	y	a	A	p	1,2,4,6
<i>Townsendia montana</i> Jones	v	a	Z	p	4,6
BORAGINACEAE					
<i>Eritrichium nanum</i> (Vill.) Schrad. var. <i>elongatum</i> (Rydb.) Cronq.	b	a	Z	p	1,4,6
<i>Mertensia ciliata</i> (James) G. Don	b	a	Z	p	2
<i>Myosotis sylvatica</i> Hoffm. var. <i>alpestris</i> (Schmidt) Koch.	b	a	Z	p	1,4
BRASSICACEAE					
<i>Arabis drummondii</i> A. Gray	v	a	Z	p	1
* <i>A. holboellii</i> Hornem.	v	a	Z	p	1
<i>A. lemmonii</i> S. Wats.	v	a	Z	p	4,6
<i>A. lyallii</i> S. Wats.	v	a	Z	p	1,6
<i>A. nuttallii</i> Robinson	w	a	Z	p	6
<i>Draba apiculata</i> Hitchc.	w	a	Z	p	1
* <i>D. aurea</i> Vahl.	y	a	Z	p	4,5,6
<i>D. crassa</i> Rydb.	y	a	Z	p	4
* <i>D. crassifolia</i> Graham	y	a	Z	a	1
<i>D. lonchocarpa</i> Rydb.	w	a	Z	p	4,5,6
<i>D. oligosperma</i> Hook.	y	a	Z	p	4
<i>D. ventosa</i> Gray	w	a	Z	p	4,5
<i>Erysimum asperum</i> (Nutt.) DC.	y	a	Z	p	6
<i>Physaria australis</i> (Pays.) Rollins	y	a	Z	p	4
* <i>Smelowskia calycina</i> C. A. Meyer var. <i>americana</i> (Regal & Herd) Drury & Rollins	w	a	Z	p	1,6
CAMPANULACEAE					
* <i>Campanula rotundifolia</i> L.	v	a	Z	p	1,4
CARYOPHYLLACEAE					
<i>Arenaria congesta</i> Nutt.	w	a	Z	p	4,6
<i>A. nuttallii</i> Pax.	w	a	Z	p	6
* <i>A. obtusiloba</i> (Rydb.) Fern	w	a	Z	p	1,4,6
* <i>Cerastium arvense</i> L.	w	a	Z	p	2
* <i>C. beeringianum</i> Cham. & Schlecht.	w	a	Z	p	4,6
* <i>Sagina saginoides</i> (L.) Karst.	w	a	Z	p	2,6
* <i>Silene acaulis</i> L.	p	a	Z	p	1,4,6
† <i>S. parryi</i> (Wats.) Hitchc. & Mag.	w	a	Z	p	6
<i>Stellaria umbellata</i> Turcz.	w	a	Z	p	2
CRASSULACEAE					
<i>Sedum debile</i> S. Wats.	y	a	Z	p	4,6
<i>Sedum lanceolatum</i> J. Torr.	y	a	Z	p	1,4,6
<i>S. rhodanthum</i> A. Gray	p	a	Z	p	2
* <i>S. rosea</i> (L.) Scop. ssp. <i>integrifolium</i> (Raf.) Hult.	v	a	Z	p	2,3
<i>S. stenopetalum</i> Pursh	y	a	Z	p	4,5

Table 1 continued.

Family	Attribute				
	1	2	3	4	5
ERICACEAE					
° <i>Arctostaphylos uva-ursi</i> (L.) Spreng.	w	a	Z	s	2,4
<i>Caultheria humifusa</i> (Grah.) Rydb.	w	a	Z	s	2
<i>Kalmia microphylla</i> (Hook.) Heller	p	a	Z	s	2,3
<i>Phyllodoce empetrifomis</i> (Sw.) D. Don	p	a	Z	s	2
<i>P. glanduliflora</i> (Hook.) Cov.	w	a	Z	s	2,4
<i>Vaccinium scoparium</i> Leib.	w	a	Z	s	6
FABACEAE					
<i>Astragalus kentrophyta</i> Gray var. <i>implexus</i> (Canby) Barneby	v	z	Z	p	1,6
° <i>Hedysarum boreale</i> Nutt.	p	z	Z	p	6
<i>H. occidentale</i> Greene	v	z	Z	p	1,4
° <i>Oxytropis campestris</i> (L.) DC.	w	z	Z	p	1,4
<i>O. deflexa</i> (Pall.) DC. var. <i>foliosa</i> Hook. Barneby	v	z	Z	p	1,4,6
GENTIANACEAE					
<i>Gentiana calycosa</i> Griseb.	b	a	Z	p	2,6
GROSSULARIACEAE					
<i>Ribes montigenum</i> McClatchie	p	a	Z	s	2,6
HYDROPHYLLACEAE					
<i>Phacelia hastata</i> Dougl. var. <i>alpina</i> (Rydb.) Cronq.	v	a	Z	p	4,6
<i>P. sericea</i> (Grah.) Gray	v	a	Z	p	1,4,6
HYPERICACEAE					
† <i>Hypericum formosum</i> H.B.K. var. <i>nortoniae</i> (Jones) Hitchc.	y	a	Z	p	6
LINACEAE					
° <i>Linum perenne</i> L. var. <i>lewisii</i> (Pursh) Eat. & Wright	b	a	Z	p	1
ONAGRACEAE					
° <i>Epilobium alpinum</i> L.	p	a	Z	p	1,4,6
° <i>E. latifolium</i> L.	p	a	Z	p	2,4,6
POLEMONIACEAE					
<i>Linanthus nuttallii</i> Gray	w	a	Z	p	1,4
<i>Phlox pulvinata</i> (Wherry) Cronq.	w	a	Z	p	4
° <i>Polemonium pulcherrimum</i> Hook.	b	a	Z	p	4
<i>P. viscosum</i> Nutt.	b	a	Z	p	4,6
POLYGONACEAE					
<i>Eriogonum ovalifolium</i> Nutt. var. <i>depressum</i> Blank.	w	a	Z	p	4,6
† <i>E. umbellatum</i> Torr. var. <i>subalpinum</i> (Greene) Jones	y	a	Z	p	4
° <i>Oxyria digyna</i> (L.) Hill	g	a	A	p	2,4,5,6
<i>Polygonum bistortoides</i> Pursh	w	a	Z	p	2,6
° <i>P. viviparum</i> L.	w	a	A	p	2
PORTULACAEAE					
<i>Claytonia lanceolata</i> Pursh	p	a	Z	p	2
<i>C. megarhiza</i> (Gray) Parry	p	a	Z	p	4
<i>Lewisia pygmaea</i> (Gray) Robins.	p	a	Z	p	4,6
<i>L. triphylla</i> (Wats.) Robins.	p	a	Z	p	4
<i>Spraguea umbellata</i> Torr.	w	a	Z	p	4
PRIMULACEAE					
° <i>Androsace septentrionalis</i> L.	w	a	Z	a	1,6
<i>Dodecatheon conjugens</i> Greene	p	a	Z	p	2
<i>D. pulchellum</i> (Raf.) Merrill	p	a	Z	p	2
<i>Primula parryi</i> A. Gray	p	a	Z	p	2,4,5

Table 1 continued.

Family	Attribute				
	1	2	3	4	5
PYROLACEAE					
<i>Pyrola dentata</i> Smith	w	a	Z	p	4
* <i>P. minor</i> L.	p	a	Z	p	2,3
RANUNCULACEAE					
<i>Anemone multifida</i> Poir. var. <i>tetonensis</i> (Porter) Hitchc.	y	a	Z	p	1,6
<i>Aquilegia flavescens</i> S. Wats.	y	a	Z	p	2,4,6
<i>Caltha leptosepala</i> DC.	w	a	Z	p	2
<i>Ranunculus eschscholtzii</i> Schlecht.	y	a	Z	p	2
var. <i>alpinus</i> (Wats.) C. L. Hitchc.					
var. <i>eschscholtzii</i>					
var. <i>suksdorfii</i> (Gray) Benson					
<i>Trollius laxus</i> Salisb.	y	a	Z	p	2
ROSACEAE					
* <i>Dryas octopetala</i> L. var. <i>angustifolia</i> C. L. Hitchc.	w	a	Z	s	4
* <i>Geum rossii</i> (R. Br.) Ser. var. <i>turbinatum</i> (Rydb.) C. L. Hitchc.	y	a	Z	p	4
<i>Ivesia gordonii</i> (Hook.) T. & G.	y	a	Z	p	1,4
<i>Potentilla brevifolia</i> Nutt. ex T. & G.	y	a	Z	p	4
<i>P. concinna</i> Rich. var. <i>rubripes</i> (Rydb.) C. L. Hitchc.	y	a	Z	p	4
<i>P. diversifolia</i> Lehm.	y	a	Z	p	1,4
<i>P. flabellifolia</i> Hook.	y	a	Z	p	2
* <i>P. fruticosa</i> L.	y	a	Z	s	1,4,6
<i>P. glandulosa</i> Lindl. var. <i>pseudorupestris</i> (Rydb.) Breit.	y	a	Z	p	6
<i>P. gracilis</i> Dougl.	y	a	Z	p	2,6
* <i>P. nivea</i> L.	y	a	Z	p	2,4,6
<i>Rubus idaeus</i> L. ssp. <i>sachalinensis</i> (Levl.) Focke	w	a	Z	p	4
* <i>Sibbaldia procumbens</i> L.	y	a	Z	s	2,4,6
RUBIACEAE					
<i>Kellogia galioides</i> Torr.	w	a	Z	p	5
SALICACEAE					
* <i>Salix arctica</i> Pall.	-	-	Z	s	1,4,6
<i>S. cascadiensis</i> Cockerell	-	-	Z	s	1,2,4
* <i>S. glauca</i> L.	-	-	Z	s	1,2
* <i>S. reticulata</i> L. ssp. <i>nivalis</i> (Hook.) Love et al.	-	-	Z	s	2
* <i>S. rotundifolia</i> L. ssp. <i>dodgeana</i> (Rydb.) Argus	-	-	Z	s	1,4
SAXIFRAGACEAE					
† <i>Mitella pentandra</i> Hook.	g	a	Z	p	2
<i>Parnassia palustris</i> L. var. <i>montanensis</i> (Fern & Rydb.) C. L. Hitchc.	w	a	Z	p	2
<i>Saxifraga adscendens</i> L.	w	a	Z	p	2,5
* <i>S. bronchialis</i> L. var. <i>austromontana</i> (Wieg.) Jones	w	a	Z	p	2,4,5,6
<i>S. debilis</i> Engelm.	w	a	Z	p	6
* <i>S. flagellaris</i> Willd.	w	a	Z	p	2,4,5
* <i>S. oppositifolia</i> L.	v	a	Z	p	2,4
<i>S. rhomboidea</i> Greene	w	a	Z	p	2,6
<i>Telesonix jamesii</i> (Torr.) Raf. var. <i>heucheriformis</i> (Rydb.) Bacigalupi	w	a	Z	p	4,5,6
SCROPHULARIACEAE					
<i>Bessya wyomingensis</i> (A. Nels.) Rydb.	v	z	Z	p	1
† <i>Castilleja miniata</i> Dougl.	r	z	Z	p	2,6
<i>C. pulchella</i> Rydb.	r	z	Z	p	1,2
<i>C. sulphurea</i> Rydb.	y	z	Z	p	1
<i>Mimulus lewisii</i> Pursh	p	z	Z	p	2,6

Table 1 continued.

Family	Attribute				
	1	2	3	4	5
<i>Pedicularis bractcosa</i> Benth.	y	z	Z	p	4
<i>P. contorta</i> Benth.	w	z	Z	p	1
° <i>P. groenlandica</i> Retz.	v	z	Z	p	2,3
<i>P. parryi</i> Gray var. <i>purpurea</i> Parry	w	z	Z	p	1
† <i>Penstemon attenuatus</i> Dougl. ssp. <i>pseudoprocerus</i> (Rydb.) Keck.	b	z	Z	p	6
<i>P. montanus</i> Greene	v	z	Z	p	4
<i>P. whippleanus</i> Gray	y	z	Z	p	1,4,6
° <i>Veronica wormskjoldii</i> Roem. & Schult.	v	z	Z	p	2,6
VALERIANACEAE					
<i>Valeriana acutiloba</i> Rydb.	w	a	Z	p	1,4
VIOLACEAE					
<i>Viola adunca</i> J. E. Smith	v	z	Z	p	2
Monocots					
CYPERACEAE					
<i>Carex albonigra</i> MacKenz.	-	-	A	g	4,6
° <i>C. atrata</i> L.	-	-	A	g	1,2
† <i>C. douglasii</i> Boott.	-	-	A	g	6
<i>C. elynoides</i> Holm	-	-	A	g	1,4,6
† <i>C. geyeri</i> Boott.	-	-	A	g	1
<i>C. haydeniana</i> Olney	-	-	A	g	1,4,6
† <i>C. illota</i> Bailey	-	-	A	g	2,3
<i>C. luzulina</i> Olney	-	-	A	g	6
<i>C. microptera</i> Mack.	-	-	A	g	1,4
<i>C. nardina</i> Fries	-	-	A	g	1,2
<i>C. nigricans</i> C. A. Mey.	-	-	A	g	1,2
<i>C. nova</i> L. Bailey	-	-	A	g	2,6
<i>C. paysonis</i> Clokey	-	-	A	g	6
<i>C. phacocephala</i> Piper	-	-	A	g	4,6
<i>C. pyrenaica</i> Wahl.	-	-	A	g	1,2,4
† <i>C. rossii</i> Boott.	-	-	A	g	1,2
JUNCACEAE					
<i>Juncus drummondii</i> E. Bey.	-	-	A	g	2,6
<i>J. mertensianus</i> Bong.	-	-	A	g	2,3,6
<i>J. parryi</i> Engelm.	-	-	A	g	1,2,4
<i>Luzula piperi</i> (Cov.) Jones	-	-	A	g	2,3,6
° <i>L. spicata</i> (L.) DC.	-	-	A	g	2,4,6
LILIACEAE					
<i>Erythronium grandiflorum</i> Pursh	y	a	Z	p	2
° <i>Lloydia scrotina</i> (L.) Sweet.	w	a	Z	p	1,4
° <i>Zigadenus elegans</i> Pursh	w	a	Z	p	2
<i>Tofieldia glutinosa</i> (Michx.) Pers.	w	a	Z	p	2
POACEAE					
<i>Agropyron caninum</i> (L.) Beauv. var. <i>latiglume</i> (Scribn. & Smith) Hitchc.	-	-	A	g	1,4,6
var. <i>andinum</i> (Scribn. & Smith) Hitchc.	-	-	A	g	1,4,6
<i>A. scribneri</i> Vasey	-	-	A	g	2
<i>Agrostis lumilis</i> Vasey	-	-	A	g	6
† <i>A. idahoensis</i> Nash	-	-	A	g	6
° <i>A. scabra</i> Willd.	-	-	A	g	1,2
<i>A. thurberiana</i> Hitchc.	-	-	A	g	2,3
<i>A. variabilis</i> Rydb.	-	-	A	g	1,4
° <i>Calamagrostis purpurascens</i> R. Br.	-	-	A	g	4,6

Table 1 continued.

Family	Attribute				
	1	2	3	4	5
<i>Danthonia intermedia</i> Vasey	-	-	A	g	1,4
<i>Deschampsia atropurpurea</i> (Wahl.) Scheele	-	-	A	gg	2
* <i>D. cespitosa</i> (L.) Beauv.	-	-	A	g	2
* <i>Festuca ovina</i> L.	-	-	A	gg	1,6
<i>Leucopoa kingii</i> (Wats.) Weber	-	-	A	g	1,6
* <i>Phleum alpinum</i> L.	-	-	A	gg	2,3,4,6
<i>Poa alpina</i> L.	-	-	A	g	1,4,6
<i>P. cusickii</i> Vasey var. <i>cusickii</i> Hitchc.	-	-	A	g	4
var. <i>epilis</i> (Scribn.) Hitchc.					
<i>P. incurva</i> Scribn. & Wms.	-	-	A	g	4
<i>P. nervosa</i> (Hook.) Vasey var. <i>whcelerii</i> (Vasey) Hitchc.	-	-	A	g	1,4
<i>P. pattersonii</i> Vasey	-	-	A	gg	1,4,6
<i>P. reflexa</i> Vasey & Scribn.	-	-	A	g	2
<i>P. rupicola</i> Nash ex Rydb.	-	-	A	g	4
<i>P. sandbergii</i> Vasey	-	-	A	g	6
* <i>Trisetum spicatum</i> (L.) Richt.	-	-	A	g	1,2,4,6

species reflects their occasional appearance above timberline, generally in the glacial cirques.

The Tetons are quite high, with 55 peaks and 7 percent of the area of the park above 10,000 feet (3050 m). Despite this, probably because the range is so precipitous, extensive stretches of alpine meadow vegetation are uncommon. Nevertheless, some extensive areas of alpine meadows can be found in the northern part of the range (e.g., Moose Basin), around Schoolroom Glacier to the west of the main peaks, in Alaska Basin, and along the Skyline Trail in the southern part of the range. Taluses and other debris accumulations, on the other hand, are abundant throughout the range and are probably the most widespread and common alpine habitats.

The major habitat types in which each species is commonly found are listed in Table 1. Although many species are characteristic of a particular habitat, other species can be found in a variety of habitats. For this reason the assignment of each species to a particular habitat or habitats should be interpreted only in a broad sense. For some species, the habitat preference is only tentative. As more information becomes available on habitat preferences and vegetation ecology in the alpine zone of the Tetons, it is hoped that a more rigorous and exact classification will be produced. Table 3 presents the major habitat types in the alpine zone using physical and

vegetative characteristics. A brief discussion of each habitat and some of the characteristic species of each follows.

Lakes and streams are common in the alpine zone of the Tetons, but no aquatics have been found. Since Scott (1966) listed a collection of *Ranunculus natans* from the alpine of the Beartooth Range, this species should be sought in the alpine of the Tetons.

Bogs, where found, generally occur close to the lower limits of the alpine zone at 9500 feet (2900 m). They commonly occur in cirques and local depressions in the major canyons, especially along streams. Mosses are common, and *Carex* and *Salix* species dominate the vascular flora. Other species found in bogs include *Pedicularis groenlandica*, *Pyrola minor*, *Sedum rosea*, *Kalmia microphylla*, and *Agrostis thurberiana*.

The bog habitat usually grades into wet meadow habitat, with any boundary between the two often difficult to find. The wet meadow habitat is common at lower elevations, particularly in cirques, around lakes and seeps, and along streams. Vegetation occurring below late-lying snowbanks is also included in this habitat type. Wet meadows are usually dominated by various Ranunculaceae (especially *Caltha leptosepala*), and *Polygonum bistortoides*, *Pedicularis groenlandica*, *Veronica wormskjoldii*, *Deschampsia cespitosa*, and *Carex*, *Potentilla* and *Salix* species. Along streams *Phyllodoce empetriformis* and *P. glanduliflora* are especially common, along

with *Epilobium latifolium*, and *Carex*, *Dodecatheon*, and *Castilleja* species. Seeps generally support a rich mixture of *Mertensia ciliata*, and *Carex*, *Mimulus*, and *Saxifraga* species, with many mosses. Ground which is exposed late by late melting snow usually supports species like *Claytonia lanceolata*, *Erythronium grandiflorum*, and *Ranunculus eschscholtzii*.

Dry meadows are fairly common. Areas like Alaska Basin, parts of Moose Basin, large

stretches along the Skyline Trail in the southern Tetons, and slopes in the major cirques consist of this habitat type. At high elevations, this habitat is composed of scattered patches of plants with many bare areas between the patches. The top of Prospectors Mountain and Hurricane Pass are good examples of this type of vegetation. The lower elevation and more extensively vegetated dry meadows are dominated by species of the Asteraceae and Poaceae families. *Astragalus*

TABLE 2. Statistical summary of the alpine flora of the Teton Range. The families with more than 10 species and the genera with 5 or more species are also listed.

Attribute	Number of species	Percent of total	
LIFE FORM			
perennial/biennial herb	149	69.0	
perennial/biennial graminoid	44	20.4	
shrub	21	9.9	
annual herb or graminoid	2	1.0	
Total	216	100.0	
FLOWER COLOR			
yellow	52	33.3	
white	48	30.8	
violet	23	14.7	
pink	20	12.8	
blue	8	5.1	
red	2	1.3	
brown/green/orange	3	2.0	
Total	156	100.0	
FLOWER SHAPE			
actinomorphic	137	87.8	
zygomorphic	19	12.2	
Total	156	100.0	
POLLINATION MODE			
zoophilous	158	75.2	
anemophilous/apomictic	52	24.8	
Total	210	100.0	
ORIGIN			
alpine but not arctic	142	65.7	
arctic and alpine	56	25.9	
subalpine	18	8.4	
Total	216	100.0	
NUMBER OF FAMILIES	36	—	
NUMBER OF GENERA	111	—	
NUMBER OF SPECIES	216	—	
LARGEST FAMILIES (number of species)		LARGEST GENERA (number of species)	
Asteraceae (39)	Poaceae (23)	<i>Carex</i> (16)	<i>Agrostis</i> (5)
Cyperaceae (16)	Brassicaceae (15)	<i>Poa</i> (8)	<i>Arabis</i> (5)
Rosaceae (13)	Scrophulariaceae (13)	<i>Potentilla</i> (8)	<i>Erigeron</i> (5)
		<i>Senecio</i> (8)	<i>Salix</i> (5)
		<i>Draba</i> (7)	<i>Sedum</i> (5)
		<i>Saxifraga</i> (6)	

kentrophyta, *Carex nigricans*, and *Juncus drummondii* are also very common. At high elevations the vegetation is dominated by *Hymenoxys grandiflora*, *Smelowskia calycina*, *Oxytropis campestris*, *Lloydia serotina*, and *Eritrichium nanum*.

Debris accumulations include taluses, scree slopes, and boulder fellfields. These habitats are often physically disturbed by mass movements. Vegetation cover is usually very low. Many species have adapted to the disturbed and often xeric conditions of debris accumulations, generally by producing extensive root or rhizome systems. Common species include *Dryas octopetala*, *Oxyria digyna*, *Cryptogramma crista*, *Senecio fremontii*, *Phacelia hastata*, *Erigeron compositus*, *Senecio longifolia*, and *Epilobium alpinum*.

Neoglacial deposits include the full range of debris accumulations, plus some unique minor habitat types (Spence 1980). Characteristic species include *Oxyria digyna*, *Senecio fremontii*, *Poa pattersonii*, *Juncus drummondii*, *Carex phaeocephala*, and *Trisetum spicatum*. Floristically, Neoglacial deposits

are among the richest of the habitat types in the alpine of the Tetons.

Rock faces and cliffs are abundant in the Tetons. The most common species found in this habitat type include *Telesonix jamesii*, *Oxyria digyna*, and various species of *Saxifraga*, *Arabis*, and *Draba*.

ACKNOWLEDGMENTS

Ivan G. Palmblad and David R. Given read and criticized an early draft of the manuscript.

LITERATURE CITED

- BENEDICT, J. B. 1973. Chronology of cirque glaciation, Colorado Front Range. *Quat. Res.* 3: 584-599.
 BILLINGS, W. D. 1978. Alpine phytogeography across the Great Basin. *Great Basin Nat. Mem.* 2:105-117.
 FRYXELL, P. A. 1957. Mode of reproduction of higher plants. *Botanical Review* 23:135-233.
 HARTMAN, R., AND R. LICHVAR. 1979. Rocky Mountain Herbarium, University of Wyoming, Laramie, Wyoming 82071. Written communication to R. J. Shaw.
 HULTÉN, E. 1968. Flora of Alaska and neighboring territories. Stanford Univ. Press, Stanford.

TABLE 3. The major habitat types in the alpine zone of the Teton Range, with some physical and vegetative characteristics noted for each type.

Habitat type	Characteristics
Lake and stream habitat	standing or moving water present.
Bog habitat	water-saturated soil throughout the growing season, sometimes flooded in early season, vegetation cover usually 100 percent, with mosses common.
Wet meadow habitat	ground flat to gently sloping, soil generally moist, obvious boulders few and scattered, usually occurs near streams, lakes, seeps, bogs, or late lying snowbanks, vegetation cover generally high (often 100 percent), usually found below 10,500 ft (3200 m).
Dry meadow habitat	ground flat to gently sloping, soil generally dry, obvious boulders few and scattered, usually occurs away from lakes, streams, seeps, bogs, or late lying snowbanks, vegetation cover moderate to high at low elevations, low at high elevations.
Debris accumulation habitat	ground flat to steeply sloping, little soil, mostly accumulations of moderate- to large-sized rock fragments, vegetation cover very low, plants usually confined to crevices between rocks.
Rock face and cliff habitat	mostly bedrock, generally steep (40° +) sometimes with ephemeral streams in early season, plants usually growing on small ledges or in cracks in the rocks.
Neoglacial deposits	flat to more often steep accumulations of glacial debris, usually near existing glaciers or permanent snowbanks, vegetation cover generally low, recently formed (100-3000 years old). See Benedict 1973, Mahaney 1975.

- JOHNSON, P. L., AND W. D. BILLINGS. 1962. The alpine vegetation of the Beartooth Plateau in relation to cryopedogenic processes and patterns. *Ecol. Monogr.* 32:105-135.
- LOVE, J. D., AND J. C. REED. 1968. Creation of the Teton landscape. Grand Teton Natural History Association Press.
- MAHANAY, W. C. 1975. Soils of post-Audubon age, Teton Glacier area, Wyoming. *Arct. & Alp. Res.* 7:141-153.
- . 1980. Department of Geography, York Univ., Atkinson College, 4700 Keele Street, Downsview, Ontario, Canada M3J 2R7. Written communication to J. R. Spence.
- MERKLE, J. 1962. Ecological studies of the Amphitheater and Surprise Lakes cirque in the Teton Mountains, Wyoming. National Park Service Files, unpublished manuscript, 23 pp.
- . 1963. Ecological studies in the Holly Lake cirque of the Teton Mountains, Wyoming. National Park Service Files, unpublished manuscript, 39 pp.
- MOLDENKE, A. R. 1976. California pollination ecology and vegetation types. *Phytologia* 34:305-361.
- OSTLER, W. K., AND K. T. HARPER. 1978. Floral ecology in relation to plant species diversity in the Wasatch Mountains of Utah. *Ecology* 59:848-861.
- PERCIVAL, M. S. 1965. *Floral biology*. Pergamon Press, London.
- POJAR, J. 1974. Reproductive dynamics of four plant communities of southwestern British Columbia. *Can. J. Bot.* 52:1819-1834.
- POLUNIN, N. 1959. *Circumpolar Arctic flora*. Clarendon Press, Oxford.
- REED, J. C., AND R. E. ZARTMAN. 1973. Geochronology of precambrian rocks of the Teton Range, Wyoming. *Geol. Soc. Amer. Bull.* 84:561-582.
- REED, J. F. 1952. Vegetation of Jackson Hole Wildlife Park, Wyoming. *Amer. Midl. Natur.* 48:700-729.
- SCOTT, R. W. 1966. The alpine flora of northwestern Wyoming. Unpublished thesis. University of Wyoming, 219 pp.
- SHAW, R. J. 1958. Vascular plants of Grand Teton National Park. *Amer. Midl. Natur.* 59:146-166.
- . 1968. Vascular plants of Grand Teton National Park, Wyoming. *Sida* 4:1-56.
- . 1976. Field guide to the vascular plants of Grand Teton National Park and Teton County, Wyoming. Utah State Univ., Logan.
- SPENCE, J. R. 1980. Vegetation of subalpine and alpine moraines in the Teton Range, Grand Teton National Park, Wyoming. Unpublished thesis. Utah State Univ., 207 pp.
- SWALES, D. E. 1979. Nectaries of certain arctic and subarctic plants with notes on pollination. *Rhodora* 81:367-407.
- WARDLE, P. 1978. Origin of the New Zealand mountain flora, with special reference to trans-Tasman relationships. *New Zealand J. Bot.* 16:535-550.