



1-31-1986

**New oriental genus of leafhoppers in the tribe Coelidiini with descriptions of new species (Homoptera: Cicadellidae: Coelidiinae)**

M. W. Nielson  
*Brigham Young University*

Follow this and additional works at: <https://scholarsarchive.byu.edu/gbn>


---

**Recommended Citation**

Nielson, M. W. (1986) "New oriental genus of leafhoppers in the tribe Coelidiini with descriptions of new species (Homoptera: Cicadellidae: Coelidiinae)," *Great Basin Naturalist*: Vol. 46 : No. 1 , Article 13. Available at: <https://scholarsarchive.byu.edu/gbn/vol46/iss1/13>

This Article is brought to you for free and open access by the Western North American Naturalist Publications at BYU ScholarsArchive. It has been accepted for inclusion in Great Basin Naturalist by an authorized editor of BYU ScholarsArchive. For more information, please contact [scholarsarchive@byu.edu](mailto:scholarsarchive@byu.edu), [ellen\\_amatangelo@byu.edu](mailto:ellen_amatangelo@byu.edu).





Figs. 1-7. *Styloidia pectinata*, n. sp.: 1. Head and pronotum, dorsal view. 2. Face. 3. Male pygofer, lateral view. 4. Connective, aedeagus, and right style, dorsal view. 5. Aedeagus, lateral view. 6. Style, lateral view. 7. Plate, ventral view.

**MALE.** Pygofer in lateral view with hooked caudodorsal process (Fig. 3); aedeagus long, narrow, tubular, slightly curved in lateral view and extending beyond apex of style, without processes (Figs. 4, 5); gonopore distad of middle of shaft on lateroventral margin; style short, not reaching apex of aedeagal shaft, with 4-5 very long nearly straight spines arising from apex, spines nearly as long as basal half of style (Figs. 4, 6); plate long and narrow (Fig. 7).

**FEMALE.** Unknown.

**Holotype** (male), MALAYSIA: Malay Peninsula, Perak, 15-XI-1927, G. Kledang (BMNH).

**REMARKS:** *S. pectinata* is easily separated from *S. cristata* by the hooked caudodorsal process on the pygofer and by the long spines that arise from the apex of the style.

Additional data on the label attached to the holotype specimen include abbreviations "F.M.S." and "2650" on the top side and

"presented by Mr. E. S. Kimund, Sel: Mus: on the underside.


*Styloidia cristata*, n. sp.

Length: Male 8.80 mm.

Medium-sized, slender species. General color blackish with numerous mottled tannish markings on forewing.

Head as in *pectinata* n. sp. but with large semiglobular eyes (Fig. 8); pronotum, scutellum and forewing as in description of genus clypeus and clypellus as in *pectinata* (Fig. 9).

**MALE.** Pygofer in lateral view with small translucent, scalelike process on caudodorsal margin (Fig. 10); aedeagus long and tubular without processes (Figs. 11, 12); gonopore as in *pectinata*; style very long, extending distally beyond apex of aedeagal shaft, with two rows of uniseriate stout spines on distal half one row on dorsal margin and the other on the inner lateral margin (Figs. 11, 13); plate as in *pectinata* (Fig. 14).



Figs. 8-14. *Styloidia cristata*, n. sp.: 8, Head and pronotum, dorsal view. 9, Face. 10, Pygofer, lateral view. 11, Connective, aedeagus, and right style, dorsal view. 12, Aedeagus, lateral view. 13, Style, lateral view. 14, Plate, ventral view.

FEMALE. Unknown.

Holotype (male) MALAYSIA: Pahang Bukit bamban, 90 km WNW of Kuala Rompin, ca 50m  
-9. X. 1961, at night, K. J. Kuncheria  
BPBM).

REMARKS: This species can be distinguished from *S. pectinata* by the long style that exceeds the length of aedeagus and by the two uniseriate rows of spines on the distal half of style.

J. Linsley Gressitt, Bernice P. Bishop Museum, Honolulu (BPBM), graciously loaned the specimens on which this paper was based. Jean Stanger provided the illustrations and Dr. James P. Kramer, U. S. National Museum, Washington, D.C., and Dr. Paul W. Oman, Oregon State University, Corvallis, gave of their time to review and improve the content of the paper.

#### LITERATURE CITED

- NIELSON, M. W. 1982. A revision of the subfamily Coelidiinae (Homoptera: Cicadellidae). IV. Tribe Coelidiini. Pacific Insects Monogr. 38. 318 pp., 1,104 figs.

#### ACKNOWLEDGMENTS

Dr. W. J. Knight, British Museum (Natural History), London (BMNH), and the late Dr.